

THE STABLE MANIFOLD THEOREM FOR SEMILINEAR STOCHASTIC EVOLUTION EQUATIONS AND STOCHASTIC PARTIAL DIFFERENTIAL EQUATIONS

II: EXISTENCE OF STABLE AND UNSTABLE MANIFOLDS

SALAH-ELDIN A. MOHAMMED*, TUSHENG ZHANG**, AND HUAIZHONG ZHAO***

ABSTRACT. This article is a sequel to [M.Z.Z.1] aimed at completing the characterization of the pathwise local structure of solutions of semilinear stochastic evolution equations (see's) and stochastic partial differential equations (spde's) near stationary solutions. Stationary solution are viewed as random points in the infinite-dimensional state space, and the characterization is expressed in terms of the almost sure long-time behavior of trajectories of the equation in relation to the stationary solution. More specifically, we establish *local stable manifold theorems* for semilinear see's and spde's (Theorems 4.1-4.4). These results give smooth stable and unstable manifolds in the neighborhood of a hyperbolic stationary solution of the underlying stochastic equation. The stable and unstable manifolds are stationary, live in a stationary tubular neighborhood of the stationary solution and are asymptotically invariant under the stochastic semiflow of the see/spde. The proof uses infinite-dimensional multiplicative ergodic theory techniques and interpolation arguments (Theorem 2.1).

1. Introduction. Hyperbolicity of a stationary trajectory.

In [M-Z-Z.1], we established the existence of perfect differentiable cocycles generated by mild solutions of a large class of semilinear stochastic evolution equations (see's) and stochastic partial differential equations (spde's). The present article is a continuation of the analysis in [M-Z-Z.1]. In this paper we introduce the concept of a stationary trajectory for the see. Within the context of stochastic differential equations (with memory) (sde's and sfde's), this concept has been used extensively in previous work of one of the authors with M. Scheutzow ([M-S.1], [M-S.2-4]). Our main objective is to characterize the pathwise local structure of solutions of semilinear see's and spde's near stationary solutions. We introduce the concept of *hyperbolicity* for a stationary solution of an see. Hyperbolicity is defined by the non-vanishing of the Lyapunov spectrum of the linearized cocycle. The hyperbolic structure of the stochastic semiflow leads to *local stable manifold theorems* (Theorems 4.1-4.4) for semilinear see's and spde's. For a hyperbolic stationary

*The research of this author is supported in part by NSF Grants DMS-9703852, DMS-9975462 and DMS-0203368.

**The research of this author is supported in part by EPSRC Grant GR/R91144.

***The research of this author is supported in part by EPSRC Grants GR/R69518 and GR/R93582.

AMS 1991 *subject classifications*. Primary 60H10, 60H20; secondary 60H25.

Key words and phrases. Stochastic semiflow, cocycle, stochastic evolution equation (see), stochastic partial differential equation (spde), multiplicative ergodic theorem, stationary solution, hyperbolicity, local stable (unstable) manifolds.

solution of the see, this gives smooth stable and unstable manifolds in a neighborhood of the stationary solution. The stable and unstable manifolds are stationary, live in a stationary tubular neighborhood of the stationary solution and are asymptotically invariant under the stochastic semiflow. The proof of the stable manifold theorem uses infinite-dimensional multiplicative ergodic theory techniques ([Ru.1], [Ru.2]) together with interpolation and perfection arguments ([Mo.1], [M-S.4]). In particular, we will assume that the reader is familiar with the results and the techniques in Ruelle's articles [Ru.1] and [Ru.2]. Our results cover semilinear stochastic evolution equations, stochastic parabolic equations, stochastic reaction-diffusion equations, and Burgers equation with additive infinite-dimensional noise.

We recall below the definition of a cocycle in Hilbert space.

Let $(\Omega, \mathcal{F}, P)$ be a complete probability space. Suppose $\theta : \mathbf{R} \times \Omega \rightarrow \Omega$ is a group of P -preserving ergodic transformations on $(\Omega, \mathcal{F}, P)$. Denote by $\bar{\mathcal{F}}$ the P -completion of $\mathcal{F}$.

Let H be a real separable Hilbert space with norm $|\cdot|$ and Borel σ -algebra $\mathcal{B}(H)$.

Take k to be any non-negative integer and $\epsilon \in (0, 1]$. Recall that a $C^{k,\epsilon}$ *perfect cocycle* (U, θ) on H is a $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{B}(H) \otimes \bar{\mathcal{F}}, \mathcal{B}(H))$ -measurable random field $U : \mathbf{R}^+ \times H \times \Omega \rightarrow H$ with the following properties:

- (i) For each $\omega \in \Omega$, the map $\mathbf{R}^+ \times H \ni (t, x) \mapsto U(t, x, \omega) \in H$ is continuous; for fixed $(t, \omega) \in \mathbf{R}^+ \times \Omega$, the map $H \ni x \mapsto U(t, x, \omega) \in H$ is $C^{k,\epsilon}$ ($D^k U(t, x, \omega)$ is C^ϵ in x on bounded subsets of H).
- (ii) $U(t_1 + t_2, \cdot, \omega) = U(t_2, \cdot, \theta(t_1, \omega)) \circ U(t_1, \cdot, \omega)$ for all $t_1, t_2 \in \mathbf{R}^+$, all $\omega \in \Omega$.
- (iii) $U(0, x, \omega) = x$ for all $x \in H, \omega \in \Omega$.

We now introduce the concept of a *stationary point* for a cocycle (U, θ) . Stationary points play the role of stochastic equilibria for the stochastic dynamical system.

Definition 1.1.

An $\mathcal{F}$ -measurable random variable $Y : \Omega \rightarrow H$ is said to be a *stationary random point* for the cocycle (U, θ) if it satisfies the following identity:

$$U(t, Y(\omega), \omega) = Y(\theta(t, \omega)) \quad (1.1)$$

for all $(t, \omega) \in \mathbf{R}^+ \times \Omega$.

The reader may note that the above definition is an infinite-dimensional analogue of a corresponding concept of invariance that was used by one of the authors in joint work with M. Scheutzwow to give a proof of the stable manifold theorem for stochastic ordinary differential equations (Definition 3.1, [M-S.3]). Definition 1.1 essentially gives a useful realization of the idea of an invariant measure for a stochastic dynamical system generated by an spde or a see. Such a realization allows us to analyze the local *almost sure* stability properties of the stochastic semiflow in the neighborhood of the stationary point. The existence (and uniqueness/ergodicity) of a stationary random point for various classes of spde's and see's has been studied by many researchers. In this article,

we will move beyond the issue of existence of stationary solutions, and apply our stable/unstable manifold theorem to examine further the almost sure asymptotic structure of the stochastic flow generated by several well-known classes of see's and spde's. In particular, we establish the existence of local stable and unstable manifolds near their stationary points.

The main objective of this section is to define the concept of *hyperbolicity* for a stationary point Y of the cocycle (U, θ) .

First, we linearize the $C^{k,\epsilon}$ cocycle (U, θ) along a stationary random point Y . By taking Fréchet derivatives at $Y(\omega)$ on each side of the cocycle identity (ii) above, using the chain rule and the definition of Y , we immediately see that $(DU(t, Y(\omega), \omega), \theta(t, \omega))$ is an $L(H)$ -valued perfect cocycle. Secondly, we appeal to the following classical result which goes back to Oseledec in the finite-dimensional case, and to D. Ruelle in infinite dimensions.

Theorem 1.1. (*Oseledec-Ruelle*)

Let $T : \mathbf{R}^+ \times \Omega \rightarrow L(H)$ be strongly measurable, such that (T, θ) is an $L(H)$ -valued cocycle, with each $T(t, \omega)$ compact. Suppose that

$$E \sup_{0 \leq t \leq 1} \log^+ \|T(t, \cdot)\|_{L(H)} + E \sup_{0 \leq t \leq 1} \log^+ \|T(1-t, \theta(t, \cdot))\|_{L(H)} < \infty.$$

Then there is a sure event $\Omega_0 \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_0) \subseteq \Omega_0$ for all $t \in \mathbf{R}^+$, and for each $\omega \in \Omega_0$, the limit

$$\Lambda(\omega) := \lim_{t \rightarrow \infty} [T(t, \omega)^* \circ T(t, \omega)]^{1/(2t)}$$

exists in the uniform operator norm. Each linear operator $\Lambda(\omega)$ is compact, non-negative and self-adjoint with a discrete spectrum

$$e^{\lambda_1} > e^{\lambda_2} > e^{\lambda_3} > \dots$$

where the λ_i 's are distinct and non-random. Each eigenvalue $e^{\lambda_i} > 0$ has a fixed finite non-random multiplicity m_i and a corresponding eigen-space $F_i(\omega)$, with $m_i := \dim F_i(\omega)$. Set $i = \infty$ when $\lambda_i = -\infty$. Define

$$E_1(\omega) := H, \quad E_i(\omega) := [\oplus_{j=1}^{i-1} F_j(\omega)]^\perp, \quad i > 1, \quad E_\infty := \ker \Lambda(\omega).$$

Then

$$E_\infty \subset \dots \subset E_{i+1}(\omega) \subset E_i(\omega) \subset \dots \subset E_2(\omega) \subset E_1(\omega) = H,$$


$$\lim_{t \rightarrow \infty} \frac{1}{t} \log |T(t, \omega)x| = \begin{cases} \lambda_i & \text{if } x \in E_i(\omega) \setminus E_{i+1}(\omega), \\ -\infty & \text{if } x \in E_\infty(\omega), \end{cases}$$

and

$$T(t, \omega)(E_i(\omega)) \subseteq E_i(\theta(t, \omega))$$

for all $t \geq 0$, $i \geq 1$.

The following figure illustrates the Oseledec-Ruelle theorem.



Proof of Theorem 1.1.

The proof is based on a discrete version of Oseledec's multiplicative ergodic theorem and the perfect ergodic theorem ([Ru.1], I.H.E.S Publications, 1979, pp. 303-304; cf. [O], [Mo.1], Lemma 5. See also Lemma 3.1 (ii) of this article). Details of the extension to continuous time are given in [Mo.1] within the context of linear stochastic functional differential equations. The arguments in [Mo.1] extend directly to general linear cocycles in Hilbert space. Cf. [F-S]. $\square$

Definition 1.2.

The sequence $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$ in the Oseledec-Ruelle theorem (Theorem 1.1) is called the *Lyapunov spectrum* of the linear cocycle (T, θ) .

Hyperbolicity of a stationary point $Y : \Omega \rightarrow H$ of the non-linear cocycle (U, θ) may now be defined in terms of a spectral gap in the Lyapunov spectrum of the linearized cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$.

Definition 1.3.

Let (U, θ) be a $C^{k,\epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) perfect cocycle on a separable Hilbert space H such that $U(t, \cdot, \omega) : H \rightarrow H$ takes bounded sets into relatively compact sets for each $(t, \omega) \in (0, \infty) \times \Omega$. A stationary point $Y(\omega)$ of the cocycle (U, θ) is *hyperbolic* if

(a) For any $a \in (0, \infty)$,

$$\int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \|DU(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega))\|_{L(H)} dP(\omega) < \infty.$$

(b) The linearized cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$ has a non-vanishing Lyapunov spectrum $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$, viz. $\lambda_i \neq 0$ for all $i \geq 1$.

By the Oseledec theorem (Theorem 1.1), the integrability condition in Definition 1.2 (a) implies the existence of a discrete Lyapunov spectrum for the linearized cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$ in Definition 1.2 (b) above.

The following result is a random version of the saddle point property for hyperbolic linear cocycles. A proof is given in ([Mo.1], Theorem 4, Corollary 2; [M-S.1], Theorem 5.3) within the context of stochastic differential systems with memory; but the arguments therein extend immediately to linear cocycles in Hilbert space.

Theorem 1.2. *(Stable and unstable subspaces)*

Let (T, θ) be a linear cocycle on a Hilbert space H . Assume that $T(t, \omega) : H \rightarrow H$ is a compact linear operator for each $t > 0$ and a.a. $\omega \in \Omega$. Suppose that

$$E \log^+ \sup_{0 \leq t_1, t_2 \leq 1} \|T(t_2, \theta(t_1, \cdot))\|_{L(H)} < \infty,$$

and let the cocycle (T, θ) have a non-vanishing Lyapunov spectrum $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$. Pick $i_0 > 1$ such that $\lambda_{i_0} < 0 < \lambda_{i_0-1}$.

Then there is a sure event $\Omega^* \in \mathcal{F}$ and stable and unstable subspaces $\{S(\omega), U(\omega) : \omega \in \Omega^*\}$, $\mathcal{F}$ -measurable (into the Grassmanian), such that for each $\omega \in \Omega^*$, the following is true:

(i) $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$.

(ii) $H = \mathcal{U}(\omega) \oplus \mathcal{S}(\omega)$. The subspace $\mathcal{U}(\omega)$ is finite-dimensional with a fixed non-random dimension, and $\mathcal{S}(\omega)$ is closed with a finite non-random codimension.

(iii) (Invariance)

$$T(t, \omega)(\mathcal{U}(\omega)) = \mathcal{U}(\theta(t, \omega)), \quad T(t, \omega)(\mathcal{S}(\omega)) \subseteq \mathcal{S}(\theta(t, \omega)),$$


for all $t \geq 0$,

(iv) (Exponential dichotomies)

$$|T(t, \omega)(x)| \geq |x|e^{\delta_1 t} \quad \text{for all } t \geq \tau_1^*, x \in \mathcal{U}(\omega),$$

$$|T(t, \omega)(x)| \leq |x|e^{-\delta_2 t} \quad \text{for all } t \geq \tau_2^*, x \in \mathcal{S}(\omega),$$

where $\tau_i^* = \tau_i^*(x, \omega) > 0, i = 1, 2$, are random times and $\delta_i > 0, i = 1, 2$, are fixed.



2. The non-linear ergodic theorem.

The main objective of this section is to refine and extend discrete-time results of D. Ruelle to the continuous-time setting in Theorem 2.1 below. This setting underlies the dynamics of the semilinear see's and spde's studied by the authors in [M-Z-Z.1]. As will be apparent later, the extension of Ruelle's results to continuous-time is non-trivial. Indeed, Section 3 in its entirety is devoted to the proof of Theorem 2.1. The main difficulties in the analysis are outlined after the statement of the theorem.

In the following, denote by $B(x, \rho)$ the open ball, radius ρ and center $x \in H$, and by $\bar{B}(x, \rho)$ the corresponding closed ball.

Theorem 2.1. (*The local stable manifold theorem*)

Let (U, θ) be a $C^{k, \epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) perfect cocycle on a separable Hilbert space H such that for each $(t, \omega) \in (0, \infty) \times \Omega$, $U(t, \cdot, \omega) : H \rightarrow H$ takes bounded sets into relatively compact sets. For any $\rho \in (0, \infty)$, denote by $\|\cdot\|_{k, \epsilon}$ the $C^{k, \epsilon}$ -norm on the space $C^{k, \epsilon}(\bar{B}(0, \rho), H)$. Let Y be a hyperbolic stationary point of the cocycle (U, θ) satisfying the following integrability property:

$$\int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \|U(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega))\|_{k, \epsilon} dP(\omega) < \infty$$

for any fixed $0 < \rho, a < \infty$ and $\epsilon \in (0, 1]$. Denote by $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$ the Lyapunov spectrum of the linearized cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega), t \geq 0)$. Define $\lambda_{i_0} := \max\{\lambda_i : \lambda_i < 0\}$ if at least one $\lambda_i < 0$. If all finite λ_i are positive, set $\lambda_{i_0} := -\infty$. (Thus λ_{i_0-1} is the smallest positive Lyapunov exponent of the linearized cocycle, if at least one $\lambda_i > 0$; in case all the λ_i 's are negative, set $\lambda_{i_0-1} := \infty$.)

Fix $\epsilon_1 \in (0, -\lambda_{i_0})$ and $\epsilon_2 \in (0, \lambda_{i_0-1})$. Then there exist

- (i) a sure event $\Omega^* \in \mathcal{F}$ with $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$,
- (ii) $\bar{\mathcal{F}}$ -measurable random variables $\rho_i, \beta_i : \Omega^* \rightarrow (0, 1)$, $\beta_i > \rho_i > 0$, $i = 1, 2$, such that for each $\omega \in \Omega^*$, the following is true:

There are $C^{k, \epsilon}$ ($\epsilon \in (0, 1]$) submanifolds $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ of $\bar{B}(Y(\omega), \rho_1(\omega))$ and $\bar{B}(Y(\omega), \rho_2(\omega))$ (resp.) with the following properties:

- (a) For $\lambda_{i_0} > -\infty$, $\tilde{\mathcal{S}}(\omega)$ is the set of all $x \in \bar{B}(Y(\omega), \rho_1(\omega))$ such that

$$|U(n, x, \omega) - Y(\theta(n, \omega))| \leq \beta_1(\omega) e^{(\lambda_{i_0} + \epsilon_1)n}$$

for all integers $n \geq 0$. If $\lambda_{i_0} = -\infty$, then $\tilde{\mathcal{S}}(\omega)$ is the set of all $x \in \bar{B}(Y(\omega), \rho_1(\omega))$ such that

$$|U(n, x, \omega) - Y(\theta(n, \omega))| \leq \beta_1(\omega) e^{\lambda n}$$

for all integers $n \geq 0$ and any $\lambda \in (-\infty, 0)$. Furthermore,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |U(t, x, \omega) - Y(\theta(t, \omega))| \leq \lambda_{i_0} \quad (2.1)$$

for all $x \in \tilde{\mathcal{S}}(\omega)$. Each stable subspace $\mathcal{S}(\omega)$ of the linearized cocycle

$(DU(t, Y(\cdot), \cdot), \theta(t, \cdot))$ is tangent at $Y(\omega)$ to the submanifold $\tilde{\mathcal{S}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = \mathcal{S}(\omega)$.

In particular, $\text{codim } \tilde{\mathcal{S}}(\omega) = \text{codim } \mathcal{S}(\omega)$, is fixed and finite.

- (b) $\limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup \left\{ \frac{|U(t, x_1, \omega) - U(t, x_2, \omega)|}{|x_1 - x_2|} : x_1 \neq x_2, x_1, x_2 \in \tilde{\mathcal{S}}(\omega) \right\} \right] \leq \lambda_{i_0}$.

(c) (Cocycle-invariance of the stable manifolds):

There exists $\tau_1(\omega) \geq 0$ such that

$$U(t, \cdot, \omega)(\tilde{\mathcal{S}}(\omega)) \subseteq \tilde{\mathcal{S}}(\theta(t, \omega)) \quad (2.2)$$

for all $t \geq \tau_1(\omega)$. Also

$$DU(t, Y(\omega), \omega)(\mathcal{S}(\omega)) \subseteq \mathcal{S}(\theta(t, \omega)), \quad t \geq 0. \quad (2.3)$$

(d) For $\lambda_{i_0-1} < \infty$, $\tilde{\mathcal{U}}(\omega)$ is the set of all $x \in \bar{B}(Y(\omega), \rho_2(\omega))$ with the property that there is a discrete-time “history” process $y(\cdot, \omega) : \{-n : n \geq 0\} \rightarrow H$ such that $y(0, \omega) = x$ and for each integer $n \geq 1$, one has $U(1, y(-n, \omega), \theta(-n, \omega)) = y(-(n-1), \omega)$ and

$$|y(-n, \omega) - Y(\theta(-n, \omega))| \leq \beta_2(\omega)e^{-(\lambda_{i_0-1} - \epsilon_2)n}.$$

If $\lambda_{i_0-1} = \infty$, $\tilde{\mathcal{U}}(\omega)$ is the set of all $x \in \bar{B}(Y(\omega), \rho_2(\omega))$ with the property that there is a discrete-time “history” process $y(\cdot, \omega) : \{-n : n \geq 0\} \rightarrow H$ such that $y(0, \omega) = x$ and for each integer $n \geq 1$,

$$|y(-n, \omega) - Y(\theta(-n, \omega))| \leq \beta_2(\omega)e^{-\lambda n},$$

for any $\lambda \in (0, \infty)$. Furthermore, for each $x \in \tilde{\mathcal{U}}(\omega)$, there is a unique continuous-time “history” process also denoted by $y(\cdot, \omega) : (-\infty, 0] \rightarrow H$ such that $y(0, \omega) = x$, $U(t, y(s, \omega), \theta(s, \omega)) = y(t+s, \omega)$ for all $s \leq 0, 0 \leq t \leq -s$, and

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |y(-t, \omega) - Y(\theta(-t, \omega))| \leq -\lambda_{i_0-1}.$$

Each unstable subspace $\mathcal{U}(\omega)$ of the linearized cocycle $(DU(t, Y(\cdot), \cdot), \theta(t, \cdot))$ is tangent at $Y(\omega)$ to $\tilde{\mathcal{U}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) = \mathcal{U}(\omega)$. In particular, $\dim \tilde{\mathcal{U}}(\omega)$ is finite and non-random.

(e) Let $y(\cdot, x_i, \omega), i = 1, 2$, be the history processes associated with

$x_i = y(0, x_i, \omega) \in \tilde{\mathcal{U}}(\omega), i = 1, 2$. Then

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup \left\{ \frac{|y(-t, x_1, \omega) - y(-t, x_2, \omega)|}{|x_1 - x_2|} : x_1 \neq x_2, x_i \in \tilde{\mathcal{U}}(\omega), i = 1, 2 \right\} \right] \\ \leq -\lambda_{i_0-1}. \end{aligned}$$

(f) (Cocycle-invariance of the unstable manifolds):

There exists $\tau_2(\omega) \geq 0$ such that

$$\tilde{\mathcal{U}}(\omega) \subseteq U(t, \cdot, \theta(-t, \omega))(\tilde{\mathcal{U}}(\theta(-t, \omega))) \quad (2.4)$$

for all $t \geq \tau_2(\omega)$. Also

$$DU(t, \cdot, \theta(-t, \omega))(\mathcal{U}(\theta(-t, \omega))) = \mathcal{U}(\omega), \quad t \geq 0;$$

and the restriction

$$DU(t, \cdot, \theta(-t, \omega))|_{\mathcal{U}(\theta(-t, \omega))} : \mathcal{U}(\theta(-t, \omega)) \rightarrow \mathcal{U}(\omega), \quad t \geq 0,$$

is a linear homeomorphism onto.


(g) The submanifolds $\tilde{\mathcal{U}}(\omega)$ and $\tilde{\mathcal{S}}(\omega)$ are transversal, viz.

$$H = T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) \oplus T_{Y(\omega)}\tilde{\mathcal{S}}(\omega).$$

Assume, in addition, that the cocycle (U, θ) is C^∞ . Then the local stable and unstable manifolds $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ are also C^∞ .

The figure below summarizes the essential features of the stable manifold theorem:

The Stable Manifold Theorem



Before we give a detailed proof of Theorem 2.1, we will outline below its basic ingredients.

An outline of the proof of Theorem 2.1:

- Since Y is a hyperbolic stationary point of the cocycle (U, θ) (Definition 1.2), then the linearized cocycle satisfies the hypotheses of “perfect versions” of the ergodic theorem and Kingman’s subadditive ergodic theorem (Lemma 3.1 (ii), (iii) in Section 3). These refined versions of the ergodic theorems give invariance of the Oseledec spaces under the *continuous-time* linearized cocycle (Theorem 1.2). Thus the stable/unstable subspaces will serve as tangent spaces to the local stable/unstable manifolds of the non-linear cocycle (U, θ) .
- Define the auxiliary perfect cocycle (Z, θ) by

$$Z(t, \cdot, \omega) := U(t, (\cdot) + Y(\omega), \omega) - Y(\theta(t, \omega)), \quad t \in \mathbf{R}^+, \omega \in \Omega.$$

This gives a “centering” of the cocycle around the stationary trajectory $Y(\theta(t))$, with the property that Z has a fixed point at $0 \in H$. Employing the continuous-time integrability estimate in Theorem 2.1, the perfect ergodic theorem and the perfect subadditive ergodic theorem, the analysis in ([Ru.2], Theorems 5.1 and 6.1) may be extended to obtain local stable/unstable manifolds for the discrete cocycle $(Z(n, \cdot, \omega), \theta(n, \omega))$ near 0. These manifolds are random objects defined for all ω which are sampled from a $\theta(t, \cdot)$ -invariant *sure event* in Ω . The translates of these manifolds by the stationary point $Y(\omega)$ correspond to local stable/unstable manifolds for $U(n, \cdot, \omega)$ near $Y(\omega)$. We then interpolate between discrete times and extend the arguments in [Ru.2] further in order to conclude that the above manifolds for the discrete-time cocycle $(U(n, \cdot, \omega), \theta(n, \omega)), n \geq 1$, also serve as local stable/unstable manifolds for the *continuous-time* cocycle (U, θ) near Y .

- It turns out that the local stable/unstable manifolds are asymptotically invariant under the continuous-time cocycle (U, θ) . For the stable manifolds, the invariance follows by arguments based on (a) a refined version of the perfect subadditive ergodic theorem (Lemma 3.2, Section 3), and (b) difficult estimates using the integrability property of Theorem 2.1 and arguments behind the proofs of Ruelle’s Theorems 4.1, 5.1 ([Ru.2]). To establish asymptotic invariance of the local unstable manifolds, we introduce the concept of a *stochastic history*

process for U , which compensates for the lack of invertibility of the cocycle. Perfection arguments similar to the above give the invariance. This completes the outline of the proof of Theorem 2.1.

A full proof of Theorem 2.1 will be given in the next section. The proof is based on a discrete-time version of the theorem given in theorems 5.1, 6.1 [Ru.2]. The extension to continuous-time is done via perfection techniques and interpolation between discrete times.

3. Proof of the local stable manifold theorem.

The main objective of this section is to give a proof of Theorem 2.1. In particular, we show that the local stable/unstable manifolds for the discrete cocycle are parametrized by sure events which are invariant under the continuous-time shift $\theta(t, \cdot) : \Omega \rightarrow \Omega$. This is achieved via a number of computations based on perfection techniques. Excursions of the cocycle between discrete times are controlled by integrability hypothesis on the cocycle (U, θ) (Theorem 2.1).

“Perfect versions” of the ergodic theorem and Kingman’s subadditive ergodic theorem will be used to construct the shift-invariant sure events appearing in the statement of the local stable manifold theorem (Theorem 2.1). These results are given in Lemmas 3.1 and 3.2 below.

The following convention will be frequently used throughout the paper:

Definition 3.1.

A family of propositions $\{P(\omega) : \omega \in \Omega\}$ is said to *hold perfectly in ω* if there is a sure event $\Omega^* \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$ and $P(\omega)$ is true for every $\omega \in \Omega^*$.

Lemma 3.1.

(i) Let $\Omega_0 \in \bar{\mathcal{F}}$ be a sure event such that $\theta(t, \cdot)(\Omega_0) \subseteq \Omega_0$ for all $t \geq 0$. Then there is a sure event $\Omega_0^* \in \mathcal{F}$ such that $\Omega_0^* \subseteq \Omega_0$ and $\theta(t, \cdot)(\Omega_0^*) = \Omega_0^*$ for all $t \in \mathbf{R}$.

(ii) Let $h : \Omega \rightarrow \mathbf{R}^+$ be any function such that there exists an $\bar{\mathcal{F}}$ -measurable function $g_1 \in L^1(\Omega, \mathbf{R}^+; P)$ and a sure event $\Omega_1 \in \bar{\mathcal{F}}$ such that $\sup_{0 \leq u \leq 1} h(\theta(u, \omega)) \leq g_1(\omega)$ for all $\omega \in \Omega_1$.

Then

$$\lim_{t \rightarrow \infty} \frac{1}{t} h(\theta(t, \omega)) = 0$$

perfectly in ω .

(iii) Suppose $f : \mathbf{R}^+ \times \Omega \rightarrow \mathbf{R} \cup \{-\infty\}$ is a process such that for each $t \in \mathbf{R}^+$, $f(t, \cdot)$ is $(\bar{\mathcal{F}}, \mathcal{B}(\mathbf{R} \cup \{-\infty\}))$ -measurable and the following conditions hold:

- (a) There is an $\bar{\mathcal{F}}$ -measurable function $g_2 \in L^1(\Omega, \mathbf{R}^+; P)$ and a sure event $\tilde{\Omega}_1 \in \bar{\mathcal{F}}$ such that $\left[\sup_{0 \leq u \leq 1} f^+(u, \omega) + \sup_{0 \leq u \leq 1} f^+(1 - u, \theta(u, \omega)) \right] \leq g_2(\omega)$ for all $\omega \in \tilde{\Omega}_1$.
- (b) $f(t_1 + t_2, \omega) \leq f(t_1, \omega) + f(t_2, \theta(t_1, \omega))$ for all $t_1, t_2 \geq 0$ and **all** $\omega \in \Omega$.

Then there is a fixed (non-random) number $f^* \in \mathbf{R} \cup \{-\infty\}$ such that

$$\lim_{t \rightarrow \infty} \frac{1}{t} f(t, \omega) = f^*$$

perfectly in ω .

Proof.

Assertion (i) is established in Proposition 2.3 ([M-S.3]).

To prove assertions (ii) and (iii) of the lemma, the reader may adapt the proofs of Lemmas 5 and 7 in [Mo.1] and employ assertion (i) above. Cf. also Lemma 3.3 in [M-S.3]. $\square$

Lemma 3.2 below is used to construct the *continuous-time* shift-invariant sure events which appear in the statement of Theorem 2.1. In essence, the lemma is a continuous-time “perfect version” of Ruelle’s Corollary A.2 ([Ru.2], p. 288).

Lemma 3.2.

Assume that the process $f : \mathbf{R}^+ \times \Omega \rightarrow \mathbf{R} \cup \{-\infty\}$ is $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{F}, \mathcal{B}(\mathbf{R} \cup \{-\infty\}))$ -measurable and satisfies the following integrability and subadditivity conditions:

- (a) $\int_{\Omega} \left[\sup_{0 \leq t_1, t_2 \leq a} f^+(t_1, \theta(t_2, \omega)) \right] dP(\omega) < \infty$ for all $a \in (0, \infty)$.
- (b) $f(t_1 + t_2, \omega) \leq f(t_1, \omega) + f(t_2, \theta(t_1, \omega))$ for all $t_1, t_2 \geq 0$ and **all** $\omega \in \Omega$.

Then there exists a fixed (non-random) $f^* \in \mathbf{R} \cup \{-\infty\}$ such that the following assertions hold perfectly in ω :

$$(i) \quad \lim_{t \rightarrow \infty} \frac{1}{t} f(t, \omega) = f^*.$$

(ii) Assume $g^* \in \mathbf{R}$ is finite and such that $f^* \leq g^*$. Then for each $\epsilon > 0$, there is an $\bar{\mathcal{F}}$ -measurable function $K_\epsilon : \Omega \rightarrow [0, \infty)$ with the following properties

$$f(t-s, \theta(s, \omega)) \leq (t-s)g^* + \epsilon t + K_\epsilon(\omega), \quad 0 \leq s \leq t < \infty,$$

$$K_\epsilon(\theta(l, \omega)) \leq K_\epsilon(\omega) + \epsilon l, \quad l \in [0, \infty).$$

Proof.

Applying Lemma 3.1 (iii), it is easy to see that there is an $f^* \in \mathbf{R} \cup \{-\infty\}$ such that assertion (i) holds for all ω in a sure event $\Omega_2 \in \mathcal{F}$ with $\theta(t, \cdot)(\Omega_2) = \Omega_2$ for all $t \in \mathbf{R}$. The integrability hypotheses (a) and Lemma 3.1 (i) imply that there is a sure event $\Omega_0 \subseteq \Omega_2$ such that $\Omega_0 \in \mathcal{F}$, $\theta(t, \cdot)(\Omega_0) = \Omega_0$ for all $t \in \mathbf{R}$, and $\sup_{0 \leq t_1, t_2 \leq a} f^+(t_1, \theta(t_2, \omega)) < \infty$ for all $a \geq 0$ and all $\omega \in \Omega_0$. Let g^* be a finite number in $[f^*, \infty)$. Define the non-negative process $g : \mathbf{R}^+ \times \Omega \rightarrow \mathbf{R}^+$ by

$$g(t, \omega) := \begin{cases} \max\{f(t, \omega) - tg^*, 0\}, & t \geq 0, \omega \in \Omega_0, \\ 0 & t \geq 0, \omega \notin \Omega_0. \end{cases}$$

Then g is $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{F}, \mathcal{B}(\mathbf{R}^+))$ -measurable and satisfies conditions (a) and (b).

Now consider the non-negative process $g' : \mathbf{R}^+ \times \Omega \rightarrow \mathbf{R}^+$ defined by

$$g'(t, \omega) := \sup_{0 \leq s \leq t} [g(s, \omega) + g(t-s, \theta(s, \omega))], \quad t \geq 0, \omega \in \Omega.$$

Observe that the projection of a $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{F})$ -measurable set is $\bar{\mathcal{F}}$ -measurable ([Co], p. 281).

Therefore, g' satisfies all the hypotheses of Lemma 3.1 (iii). This gives a non-negative g'^* such that $\lim_{t \rightarrow \infty} \frac{1}{t} g'(t, \omega) = g'^*$ for all ω in a sure event $\Omega_3 \in \mathcal{F}$, with $\theta(t, \cdot)(\Omega_3) = \Omega_3$ for all $t \in \mathbf{R}$.

We will show next the following convergence in probability:

$$\lim_{t \rightarrow \infty} \frac{1}{t} \sup_{0 \leq s \leq t} g(t-s, \theta(s, \cdot)) = 0. \quad (3.1)$$

To do this, observe that the process $h : \mathbf{R}^+ \times \Omega \rightarrow \mathbf{R}$, $h(t, \omega) := g(t, \theta(-t, \omega))$, $t \in \mathbf{R}^+$, $\omega \in \Omega$, satisfies the conditions of Lemma 3.1 (iii). Therefore

$$\lim_{t \rightarrow \infty} \frac{1}{t} h(t, \cdot) = 0$$

almost surely and hence in probability. Pick $\delta, t_0 > 0$ such that $P(\frac{1}{t} h(t, \cdot) \geq \delta) < \delta$ for all $t \geq t_0$.

Let $t \geq t_0$. Then

$$\begin{aligned} \sup_{0 \leq s \leq t} \frac{1}{t} g(t-s, \theta(s, \omega)) &\leq \sup_{0 \leq s \leq t-t_0} \frac{1}{t} g(t-s, \theta(s, \omega)) + \sup_{t-t_0 \leq s \leq t} \frac{1}{t} g(t-s, \theta(s, \omega)) \\ &\leq \sup_{0 \leq s \leq t-t_0} \frac{1}{t} g(t-s, \theta(-(t-s), \theta(t, \omega))) + \sup_{t-t_0 \leq s \leq t} \frac{1}{t} g(t-s, \theta(s, \omega)). \end{aligned}$$

By condition (a), the second term in the right hand side of the last inequality converges to zero in probability. The probability that the first term is less than or equal to δ is at least $1 - \delta$. Hence (3.1) holds.

It follows easily from (3.1) that $g'^* = 0$. This implies that assertion (i) holds for all ω in a sure event $\Omega_4 \in \mathcal{F}$ with $\Omega_4 \subseteq \Omega_0 \cap \Omega_3$ and $\theta(t, \cdot)(\Omega_4) = \Omega_4$ for all $t \in \mathbf{R}$. To complete the proof of assertion (ii), let $\epsilon > 0$ and define the $(\bar{\mathcal{F}}, \mathcal{B}(\mathbf{R}^+))$ -measurable function $K_\epsilon : \Omega_4 \rightarrow [0, \infty)$ by

$$K_\epsilon(\omega) := \sup_{0 \leq s \leq t < \infty} [g(t-s, \theta(s, \omega)) - \epsilon t]$$

for all $\omega \in \Omega_4$. This completes the proof of the lemma. $\square$

Lemma 3.3 below is essentially a “perfect version” of Proposition 3.2 in [Ru.2], p. 257. Our Lemma 3.2 plays a crucial role in the proof of Lemma 3.3. In the statement of the lemma, we will use $\mathcal{B}_s(L(H))$ to denote the Borel σ -algebra on $L(H)$ generated by the strong topology on $L(H)$, viz. the smallest topology on $L(H)$ for which all evaluations $L(H) \ni A \mapsto A(z) \in H, z \in H$, are continuous.

Lemma 3.3.

Suppose $(T^t(\omega), \theta(t, \omega))$, $t \geq 0$, is a perfect cocycle of bounded linear operators in H satisfying the following hypotheses:

- (i) The process $\mathbf{R}^+ \times \Omega \ni (t, \omega) \mapsto T^t(\omega) \in L(H)$ is $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{F}, \mathcal{B}_s(L(H)))$ -measurable.
- (ii) The map $\mathbf{R}^+ \times \Omega \ni (t, \omega) \mapsto \theta(t, \omega) \in \Omega$ is $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{F}, \mathcal{F})$ -measurable, and is a group of ergodic P -preserving transformations on $(\Omega, \mathcal{F}, P)$.
- (iii) $E \sup_{0 \leq t_1, t_2 \leq a} \log^+ \|T^{t_2}(\theta(t_1, \cdot))\|_{L(H)} < \infty$ for any finite $a > 0$.
- (iv) For each $t > 0$, $T^t(\omega)$ is compact, perfectly in ω .
- (v) For any $u \in H$, the map $[0, \infty) \ni t \mapsto T^t(\omega)(u) \in H$ is continuous, perfectly in ω .

Let $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$ be the Lyapunov spectrum of $(T^t(\omega), \theta(t, \omega))$, with Oseledec spaces

$$\dots E_{i+1}(\omega) \subset E_i(\omega) \subset \dots \subset E_2(\omega) \subset E_1(\omega) = H.$$

Let $j_0 \geq 1$ be any fixed integer with $\lambda_{j_0} > -\infty$. Let the integer function $r : \{1, 2, \dots, Q\} \rightarrow \{1, 2, \dots, j_0\}$ “count” the multiplicities of the Lyapunov exponents in the sense that $r(1) = 1$, $r(Q) = j_0$, and for each $1 \leq i \leq j_0$, the number of integers in $r^{-1}(i)$ is the multiplicity of λ_i . Set $V_n(\omega) := E_{j_0+1}(\theta(n, \omega))$, $n \geq 0$.

Then the sequence $T_n(\omega) := T^1(\theta((n-1), \omega))$, $n \geq 1$, satisfies Condition (S) of ([Ru.2], pp. 256-257) perfectly in ω with $Q = \text{codim } E_{j_0+1}(\omega)$. In particular, there is an $\mathcal{F}$ -measurable set of Q orthonormal vectors $\{\xi_0^{(1)}(\omega), \dots, \xi_0^{(Q)}(\omega)\}$ such that $\xi_0^{(k)}(\omega) \in [E_{r(k)}(\omega) \setminus E_{r(k)+1}(\omega)]$ for $k = 1, \dots, Q$, perfectly in ω , and satisfying the following properties:

$$\text{Set } \xi_t^{(k)}(\omega) := \frac{T^t(\omega)(\xi_0^{(k)}(\omega))}{|T^t(\omega)(\xi_0^{(k)}(\omega))|}, \text{ and for any } u \in H, \text{ write}$$

$$u = \sum_{k=1}^Q u_t^{(k)}(\omega) \xi_t^{(k)}(\omega) + u_t^{(Q+1)}(\omega), \quad u_t^{(Q+1)}(\omega) \in V_0(\theta(t, \omega)), \quad \omega \in \Omega.$$

Then for any $\epsilon > 0$, there is an $\bar{\mathcal{F}}$ -measurable random constant $D_\epsilon(\omega) > 0$ such that the following inequalities hold perfectly in ω :

$$|u_t^{(k)}(\omega)| \leq D_\epsilon(\omega)e^{\epsilon t}|u|$$

$$|u_t^{(Q+1)}(\omega)| \leq D_\epsilon(\omega)e^{\epsilon t}|u|$$

$$D_\epsilon(\theta(l, \omega)) \leq D_\epsilon(\omega)e^{\epsilon l}$$

for all $t \geq 0$, $1 \leq k \leq Q$ and for all $l \in [0, \infty)$.

Furthermore, all the random constants in Ruelle's condition (S) ([Ru.2], pp. 256-257) may be chosen to be $\bar{\mathcal{F}}$ -measurable in ω .

Proof.

Our proof runs along similar lines to that of Proposition 3.2 in [Ru.2]: However, one has to maintain the non-trivial requirement that all relevant arguments hold perfectly in ω .

It is assumed throughout this proof that the reader is familiar with Ruelle's conditions (S): (S1)-(S4) as spelled out in ([Ru.2], pp. 256-257).

Observe first that $T_n(\omega)$ satisfies (S1) perfectly in ω . This holds because of (iii), the perfect cocycle property, Lemma 3.1 and the proof of Theorem 4 ([Mo.1]). Note that, by the ordering of the fixed Lyapunov spectrum, relation (3.4) of [Ru.2] holds perfectly. Denote by Ω^* the $\theta(t, \cdot)$ -invariant sure event where (S1) holds. Using ergodicity of θ and the fact that $\text{codim } V_0(\omega) = Q$, for all $\omega \in \Omega^*$, it follows that $\text{codim } V_n(\omega) = \text{codim } E_{j_0+1}(\theta(n, \omega)) = Q$. Therefore, (S2) is satisfied for all $\omega \in \Omega^*$.

We next prove that (S3) holds perfectly. To do this, we will prove the stronger assertion that the *continuous-time* cocycle $(T^t(\omega), \theta(t, \omega))$ satisfies (S3) perfectly in ω . Set $\hat{T}^t(\omega) := T^t(\omega)|_{V_0(\omega)}$, $\omega \in \Omega^*$, $t \geq 0$. Hence $\hat{T}^t(\omega)(V_0(\omega)) \subseteq V_0(\theta(t, \omega))$, and the following cocycle identity

$$\hat{T}^{t_1+t_2}(\omega) = \hat{T}^{t_2}(\theta(t_1, \omega)) \circ \hat{T}^{t_1}(\omega)$$

holds for all $\omega \in \Omega^*$, $t \geq 0$. Denote $F_t(\omega) := \log \|\hat{T}^t(\omega)\|$, $\omega \in \Omega^*$, $t \geq 0$. Hypothesis (iii) of the lemma easily implies that $E \sup_{0 \leq t_1, t_2 \leq a} F_{t_2}^+(\theta(t_1, \cdot)) < \infty$ for any finite $a > 0$. Furthermore,

$(F_t(\omega), \theta(t, \omega))$ is perfectly subadditive because of the above cocycle identity. Applying Lemma 3.1, we obtain a fixed number $F^* \in \mathbf{R} \cup \{-\infty\}$ such that

$$\lim_{t \rightarrow \infty} \frac{1}{t} F_t(\omega) = F^*$$

perfectly in ω . Suppose $S = j_0$. When $\lambda_{j_0+1} > -\infty$, set $\mu^{(S+1)} := \lambda_{j_0+1}$; and when $\lambda_{j_0+1} = -\infty$, take $\mu^{(S+1)}$ to be any fixed number in $(-\infty, \lambda_{j_0})$. Using (3.5), p. 257 of [Ru.2], it follows that $F^* \leq \mu^{(S+1)}$. Suppose $\epsilon > 0$ and $\lambda_{j_0+1} > -\infty$. Then by Lemma 3.2(ii), there is an $\bar{\mathcal{F}}$ -measurable function $K_\epsilon : \Omega \rightarrow [0, \infty)$ such that

$$\log \|\hat{T}^{t-s}(\theta(s, \omega))\| \leq (t-s)\mu^{(S+1)} + \epsilon t + K_\epsilon(\omega), \quad 0 \leq s \leq t < \infty, \quad (3.2)$$

and

$$K_\epsilon(\theta(l, \omega)) \leq K_\epsilon(\omega) + \epsilon l, \quad l \in [0, \infty),$$

perfectly in ω . When $\lambda_{j_0+1} = -\infty$, the inequality (3.2) holds where $\mu^{(S+1)}$ is replaced by any (finite) number in $(-\infty, \lambda_{j_0})$. Now let m, n be positive integers such that $m < n$. In (3.2), replace t by n and s by $m+1$ to see that $T_n(\omega)$, $n \geq 1$, satisfies (S3) perfectly in ω .

The rest of this proof will now focus on showing that the sequence $T_n(\omega)$, $n \geq 1$, also satisfies Ruelle's condition (S4) perfectly in ω . Indeed, we will establish the stronger statement that the continuous-time cocycle $(T^t(\omega), \theta(t, \omega))$ satisfies (S4) perfectly in ω . Using the orthogonal decomposition $H = V_0(\theta(t, \omega)) \oplus V_0(\theta(t, \omega))^\perp$, write

$$T^t(\omega)(\xi) = \tilde{T}^t(\omega)(\xi) + \bar{T}^t(\omega)(\xi), \quad \xi \in H, t \geq 0, \omega \in \Omega^*. \quad (3.3)$$

That is, $\tilde{T}^t(\omega)(\xi) \in V_0(\theta(t, \omega))$ and $\bar{T}^t(\omega)(\xi) \in V_0(\theta(t, \omega))^\perp$ are the orthogonal projections of $T^t(\omega)(\xi)$ on $V_0(\theta(t, \omega))$ and $V_0(\theta(t, \omega))^\perp$, respectively. Thus (3.3) defines a family of continuous linear operators $\tilde{T}^t(\omega) : H \rightarrow V_0(\theta(t, \omega))^\perp \subseteq H$, $\bar{T}^t(\omega) : H \rightarrow V_0(\theta(t, \omega)) \subseteq H$, $t \geq 0$. We now

show that the family $(\check{T}^t(\omega), \theta(t, \omega)), \omega \in \Omega$, satisfies the perfect cocycle property in $L(H)$. To prove this, we fix any $\omega \in \Omega$. Then by the cocycle property of $(T^t(\omega), \theta(t, \omega))$ and (3.3), we obtain

$$\begin{aligned} T^{t_1+t_2}(\omega)(\xi) &= T^{t_2}(\theta(t_1, \omega))[T^{t_1}(\omega)(\xi)] \\ &= \check{T}^{t_2}(\theta(t_1, \omega))[\check{T}^{t_1}(\omega)(\xi)] + \check{T}^{t_2}(\theta(t_1, \omega))[\tilde{T}^{t_1}(\omega)(\xi)] + \check{T}^{t_2}(\theta(t_1, \omega))[\tilde{T}^{t_1}(\omega)(\xi)] \\ &\quad + \check{T}^{t_2}(\theta(t_1, \omega))[\tilde{T}^{t_1}(\omega)(\xi)]. \end{aligned} \quad (3.4)$$

for all $t_1, t_2 \geq 0, \xi \in H$. Furthermore, $\check{T}^t(\omega)(\xi) = 0$ for all $\xi \in V_0(\omega)$, because $V_0(\omega)$ is invariant under the cocycle $(T^t(\omega), \theta(t, \omega))$. Thus, $\check{T}^{t_2}(\theta(t_1, \omega))[\tilde{T}^{t_1}(\omega)(\xi)] = 0$ for all $\xi \in H$, and (3.4) yields

$$\begin{aligned} T^{t_1+t_2}(\omega)(\xi) &= \check{T}^{t_2}(\theta(t_1, \omega))[\check{T}^{t_1}(\omega)(\xi)] + \check{T}^{t_2}(\theta(t_1, \omega))[\tilde{T}^{t_1}(\omega)(\xi)] + \check{T}^{t_2}(\theta(t_1, \omega))[\tilde{T}^{t_1}(\omega)(\xi)]. \end{aligned} \quad (3.5)$$

Now

$$T^{t_1+t_2}(\omega)(\xi) = \check{T}^{t_1+t_2}(\omega)(\xi) + \tilde{T}^{t_1+t_2}(\omega)(\xi) \quad (3.6)$$

for all $\xi \in H$. In the right hand side of (3.5), the first term belongs to $V_0(\theta(t_1 + t_2, \omega))^\perp$, while the second two terms belong to $V_0(\theta(t_1 + t_2, \omega))$. So by uniqueness of the orthogonal decomposition, it follows from (3.6) and (3.5) that

$$\check{T}^{t_1+t_2}(\omega)(\xi) = \check{T}^{t_2}(\theta(t_1, \omega))[\check{T}^{t_1}(\omega)(\xi)] \quad (3.7)$$

for all $\xi \in H$. Hence $(\check{T}^t(\omega), \theta(t, \omega))$ is a perfect cocycle in $L(H)$.

We next verify that both cocycles $(T^t(\omega), \theta(t, \omega))$ and $(\check{T}^t(\omega), \theta(t, \omega))$ satisfy the conditions of the perfect Oseledec theorem (Theorem 1.1). To see this, note that

$$E \sup_{0 \leq t_1, t_2 \leq a} \log^+ \|\check{T}^{t_2}(\theta(t_1, \cdot))\|_{L(H)} < \infty \quad (3.8)$$

for any finite $a > 0$. This follows immediately from the integrability property (iii) of the lemma.

Now apply Theorem 1.1 to $(T^t(\omega), \theta(t, \omega))$ and $(\check{T}^t(\omega), \theta(t, \omega))$. This gives the following limits

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log |\check{T}^t(\omega)(\xi)| = \check{l}_\xi, \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log |T^t(\omega)(\xi)| = l_\xi$$

perfectly in ω for all $\xi \in H$, with $l_\xi, \check{l}_\xi$ fixed numbers in $\mathbf{R} \cup \{-\infty\}$. We now apply (3.6) in ([Ru.2], p. 259) to obtain

$$\check{l}_\xi = \lim_{n \rightarrow \infty} \frac{1}{n} \log |\check{T}^n(\omega)(\xi)| = \lim_{n \rightarrow \infty} \frac{1}{n} \log |T^n(\omega)(\xi)| = l_\xi$$

for a.a. ω and all $\xi \in H \setminus V_0(\omega)$. Therefore the equality

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log |\check{T}^t(\omega)(\xi)| = \lim_{t \rightarrow \infty} \frac{1}{t} \log |T^t(\omega)(\xi)|$$

holds perfectly in ω for all $\xi \in H \setminus V_0(\omega)$. Hence, relation (3.6) in ([Ru.2], p. 259) may be replaced by the continuous-time “perfect” relation

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \frac{|\check{T}^t(\omega)(\xi)|}{|T^t(\omega)(\xi)|} = 0 \quad (3.9)$$

for all $\xi \in H \setminus V_0(\omega)$.

By ([C-V], Theorem III.6, p. 65) and Gram-Schmidt orthogonalization, we may select a set of Q , $\mathcal{F}$ -measurable, orthonormal vectors $\{\xi_0^{(1)}(\omega), \dots, \xi_0^{(Q)}(\omega)\}$ such that $\xi_0^{(k)}(\omega) \in [E_{r(k)}(\omega) \setminus E_{r(k)+1}(\omega)] \cap V_0(\omega)^\perp$ for $k = 1, \dots, Q$, perfectly in ω . In the argument in [Ru.2], p. 259, replace (3.6) by (3.9) above, n by t , $\xi_n^{(k)}$ by $\xi_t^{(k)}(\omega) := \frac{T^t(\omega)(\xi_0^{(k)}(\omega))}{|T^t(\omega)(\xi_0^{(k)}(\omega))|}$, V_n by $V_0(\theta(t, \omega))$, and $\eta_n^{(k)}$ by $\eta_t^{(k)}(\omega) := \frac{\check{T}^t(\omega)(\xi_0^{(k)}(\omega))}{|T^t(\omega)(\xi_0^{(k)}(\omega))|}$. Therefore for $u \in H$, we write

$$u = \sum_{k=1}^Q u_t^{(k)}(\omega) \xi_t^{(k)}(\omega) + u_t^{(Q+1)}(\omega), \quad u_t^{(Q+1)}(\omega) \in V_0(\theta(t, \omega)), \quad (3.10)$$

perfectly in ω for all $t \geq 0$. Furthermore, as in [Ru.2], p. 259, (3.9) implies that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log |\det(\eta_t^{(1)}(\omega), \dots, \eta_t^{(Q)}(\omega))| = 0, \quad (3.11)$$

perfectly in ω .

It remains to prove that for each $\epsilon > 0$, there is an $\bar{\mathcal{F}}$ -measurable non-negative function $D_\epsilon : \Omega \rightarrow (0, \infty)$ such that the following inequalities

$$\left. \begin{aligned} |u_t^{(k)}(\omega)| &\leq D_\epsilon(\omega) e^{\epsilon t} |u| \\ |u_t^{(Q+1)}(\omega)| &\leq D_\epsilon(\omega) e^{\epsilon t} |u| \\ D_\epsilon(\theta(t, \omega)) &\leq D_\epsilon(\omega) e^{\epsilon t} \end{aligned} \right\} \quad (3.12)$$

hold perfectly in ω , for all $t \geq 0$, $1 \leq k \leq Q$ and for all $l \in [0, \infty)$.

In order to establish the inequalities (3.12), we define

$$D_\epsilon(\omega) := 1 + Q \cdot \sup_{0 \leq s \leq t < \infty} e^{-\epsilon t} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \eta_{t-s}^{(2)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \quad (3.13)$$

perfectly in ω .

First of all we must show that $D_\epsilon(\omega)$ is finite perfectly in ω . Surprisingly, this will require some work. Let $0 \leq s \leq t$. Observe that the determinant of the linear operator $\tilde{T}^{t-s}(\theta(s, \omega))$ is given by $\frac{|\wedge_{k=1}^Q \tilde{T}^{t-s}(\theta(s, \omega))(v_k)|}{|\wedge_{k=1}^Q v_k|}$ for *any choice of basis* $\{v_1, \dots, v_Q\}$ in $V_0(\theta(s, \omega))^\perp$. Therefore, the following inequalities hold perfectly in ω :

$$\begin{aligned} & |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \\ &= \frac{\Pi_{k=1}^Q |T^{t-s}(\theta(s, \omega))(\xi_0^{(k)}(\theta(s, \omega)))|}{|\det(\tilde{T}^{t-s}(\theta(s, \omega))(\xi_0^{(1)}(\theta(s, \omega))), \dots, \tilde{T}^{t-s}(\theta(s, \omega))(\xi_0^{(Q)}(\theta(s, \omega))))|} \\ &= \frac{\Pi_{k=1}^Q [|T^{t-s}(\theta(s, \omega))(\xi_0^{(k)}(\theta(s, \omega)))|] \cdot |\wedge_{k=1}^Q [\tilde{T}^s(\omega)(\xi_0^{(k)}(\omega))]|}{|\det(\tilde{T}^{t-s}(\theta(s, \omega))(\tilde{T}^s(\omega)(\xi_0^{(1)}(\omega))), \dots, \tilde{T}^{t-s}(\theta(s, \omega))(\tilde{T}^s(\omega)(\xi_0^{(Q)}(\omega))))|} \\ &\leq \frac{\Pi_{k=1}^Q [|T^{t-s}(\theta(s, \omega))(\xi_0^{(k)}(\theta(s, \omega)))|] \cdot |\tilde{T}^s(\omega)(\xi_0^{(k)}(\omega))|]}{|\det(\tilde{T}^t(\omega)(\xi_0^{(1)}(\omega)), \dots, \tilde{T}^t(\omega)(\xi_0^{(Q)}(\omega)))|} \\ &= \frac{\Pi_{k=1}^Q [|T^{t-s}(\theta(s, \omega))(\xi_0^{(k)}(\theta(s, \omega)))|] \cdot |\tilde{T}^s(\omega)(\xi_0^{(k)}(\omega))|]}{\|[\tilde{T}^t(\omega)|V_0(\omega)^\perp]^{\wedge Q}\|} \quad (3.14) \end{aligned}$$

$$\leq \frac{\|T^{t-s}(\theta(s, \omega))\|^Q \cdot \|\tilde{T}^s(\omega)\|^Q}{\|[\tilde{T}^t(\omega)|V_0(\omega)^\perp]^{\wedge Q}\|}. \quad (3.15)$$

By the integrability condition (iii), it follows that

$$\sup_{0 \leq s \leq t \leq a} \|T^{t-s}(\theta(s, \omega))\|^Q \cdot \|\tilde{T}^s(\omega)\|^Q < \infty$$

perfectly in ω for any finite $a > 0$.

We now prove that for each finite $a > 0$,

$$\sup_{0 \leq s \leq t \leq a} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} < \infty \quad (3.16)$$

perfectly in ω . To see this, define the compact set

$$S(\omega) := \{(t, v_1, \dots, v_Q) : t \in [0, a], v_k \in V_0(\omega)^\perp, |v_k| = 1, \langle v_k, v_l \rangle = 0, 1 \leq k < l \leq Q\}.$$

for $\omega \in \Omega$. Thus (3.16) will hold if we prove that

$$\inf_{(t, v_1, \dots, v_Q) \in S(\omega)} |\wedge_{k=1}^Q [\check{T}^t(\omega)(v_k)]| > 0 \quad (3.17)$$

perfectly in ω .

To prove (3.17), we observe that each map $\check{T}^t(\omega)|V_0(\omega)^\perp : V_0(\omega)^\perp \rightarrow V_0(\theta(t, \omega))^\perp$ is injective for each $t \geq 0$ perfectly in ω . This is an easy consequence of the cocycle property and the fact that $\lambda_{j_0} > -\infty$. In fact,

$$|\wedge_{k=1}^Q [\check{T}^t(\omega)(v_k)]| > 0 \quad (3.18)$$

for all $(t, v_1, \dots, v_Q) \in S(\omega)$. Furthermore, the map

$$[0, a] \times [V_0(\omega)^\perp]^Q \ni (t, v_1, \dots, v_Q) \mapsto |\wedge_{k=1}^Q [\check{T}^t(\omega)(v_k)]| \in [0, \infty)$$

is jointly continuous, by hypothesis (v) of the lemma. By compactness of $S(\omega)$, (3.18) implies (3.17). Therefore, (3.16) follows from (3.15) and (3.17).

The following convergence

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \sup_{0 \leq s \leq t} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} = 0 \quad (3.19)$$

holds perfectly in ω . To prove this convergence, note that (3.14) implies the following estimate

$$\begin{aligned} & |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \\ & \leq \frac{\prod_{k=1}^Q \{ \| [T^{t-s}(\theta(s, \omega))|E_{r(k)}(\theta(s, \omega))] \| \cdot \| [\check{T}^s(\omega)|E_{r(k)}(\omega)] \| \}}{\| [\check{T}^t(\omega)|V_0(\omega)^\perp]^{\wedge Q} \|} \end{aligned}$$

for $0 \leq s \leq t$ perfectly in ω . Let $\epsilon > 0$ be arbitrary. Taking $\frac{1}{t} \log \sup_{0 \leq s \leq t}$ on both sides of the above inequality and applying Lemma 3.2(ii) yields the following inequalities

$$\begin{aligned} & \frac{1}{t} \log \sup_{0 \leq s \leq t} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \\ & \leq \frac{1}{t} \sup_{0 \leq s \leq t} \left\{ \sum_{k=1}^Q (\log \| [T^{t-s}(\theta(s, \omega))|E_{r(k)}(\theta(s, \omega))] \| + \log \| [\check{T}^s(\omega)|E_{r(k)}(\omega)] \|) \right\} \\ & \quad - \frac{1}{t} \log \| [\check{T}^t(\omega)|V_0(\omega)^\perp]^{\wedge Q} \| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{t} \sup_{0 \leq s \leq t} \left\{ \sum_{k=1}^Q (t-s) \lambda_{r(k)} + \epsilon t + K_\epsilon^1(\omega) + \sum_{k=1}^Q s \lambda_{r(k)} + \epsilon s + K_\epsilon^2(\omega) \right\} \\
&\quad - \frac{1}{t} \log \| [\check{T}^t(\omega) | V_0(\omega)^\perp]^{\wedge Q} \| \\
&= \sum_{k=1}^Q \lambda_{r(k)} + 2\epsilon + \frac{1}{t} [K_\epsilon^1(\omega) + K_\epsilon^2(\omega)] - \frac{1}{t} \log \| [\check{T}^t(\omega) | V_0(\omega)^\perp]^{\wedge Q} \|, \quad t > 0,
\end{aligned}$$

perfectly in ω , with $K_\epsilon^i(\omega)$, $i = 1, 2$, finite positive random constants independent of t . Therefore, the above inequality implies that

$$\begin{aligned}
\limsup_{t \rightarrow \infty} \frac{1}{t} \log \sup_{0 \leq s \leq t} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \\
\leq \sum_{k=1}^Q \lambda_{r(k)} + 2\epsilon - \liminf_{t \rightarrow \infty} \frac{1}{t} \log \| [\check{T}^t(\omega) | V_0(\omega)^\perp]^{\wedge Q} \| \\
= \sum_{k=1}^Q \lambda_{r(k)} + 2\epsilon - \sum_{k=1}^Q \lambda_{r(k)} \\
= 2\epsilon.
\end{aligned}$$

Since $\epsilon > 0$ is arbitrary, then

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \sup_{0 \leq s \leq t} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \leq 0 \quad (3.20)$$

perfectly in ω . The convergence (3.11) immediately implies the inequality

$$\begin{aligned}
\liminf_{t \rightarrow \infty} \frac{1}{t} \log \sup_{0 \leq s \leq t} |\det(\eta_{t-s}^{(1)}(\theta(s, \omega)), \dots, \eta_{t-s}^{(Q)}(\theta(s, \omega)))|^{-1} \\
\geq \liminf_{t \rightarrow \infty} \frac{1}{t} \log |\det(\eta_t^{(1)}(\omega), \dots, \eta_t^{(Q)}(\omega))|^{-1} = 0 \quad (3.21)
\end{aligned}$$

Thus (3.19) follows from (3.20) and (3.21).

From (3.16), (3.19) and (3.13), we conclude that $D_\epsilon(\omega)$ is finite perfectly in ω .

From the definition (3.13) of $D(\omega)$, one immediately gets the last inequality in (3.12).

It remains to show the first two inequalities in (3.12). In the right hand side of (3.10), we look at the terms

$$\check{u}(\omega) = \sum_{k=1}^Q u_t^{(k)}(\omega) \eta_t^{(k)}(\omega), \quad u \in H, t \geq 0,$$

where $\check{u}(\omega), \eta_t^{(k)}(\omega), 1 \leq k \leq Q$, are viewed as column vectors in $\mathbf{R}^Q$ with respect to the basis $\{\xi_0^{(k)}(\theta(t, \omega)) : 1 \leq k \leq Q\}$. Using Cramer's rule, the above equation may be solved for each $u_t^{(k)}(\omega)$. In view of (3.13), this yields the following estimates

$$\begin{aligned}
|u_t^{(k)}(\omega)| &= \left| \frac{\det(\eta_t^{(1)}(\omega), \dots, \eta_t^{(k-1)}(\omega), \check{u}(\omega), \eta_t^{(k+1)}(\omega), \dots, \eta_t^{(Q)}(\omega))}{\det(\eta_t^{(1)}(\omega), \dots, \eta_t^{(Q)}(\omega))} \right| \\
&\leq \frac{|\check{u}(\omega)|}{|\det(\eta_t^{(1)}(\omega), \dots, \eta_t^{(Q)}(\omega))|} \\
&\leq \frac{[D_\epsilon(\omega) - 1]}{Q} |u| e^{\epsilon t} \\
&\leq D_\epsilon(\omega) |u| e^{\epsilon t}, \quad 1 \leq k \leq Q, t \geq 0,
\end{aligned} \tag{3.22}$$

perfectly in ω . By virtue of (3.10), the triangle inequality and (3.22), one gets

$$|u_t^{(Q+1)}(\omega)| \leq |u| + \sum_{k=1}^Q |u_t^{(k)}(\omega)| \leq D_\epsilon(\omega) |u| e^{\epsilon t}, \quad t \geq 0,$$

perfectly in ω . Therefore, $T_n(\omega)$ satisfies (S4) perfectly in ω , and the proof of the proposition is now complete. $\square$

In Lemma 3.4 below, an integrability condition allows us to pass from discrete-time limits of the cocycle to continuous ones. This property is crucial to the proof of Theorem 2.1. The reason this property holds is because the integrability hypothesis together with the perfect ergodic theorem (Lemma 3.1 (ii)) allow for control of the excursions of the continuous-time cocycle between discrete times.

Lemma 3.4.

Let $Y : \Omega \rightarrow H$ be a stationary point of the cocycle (U, θ) satisfying the integrability condition

$$\int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \|U(t_2, Y(\theta(t_1, \omega)) + (\cdot), \theta(t_1, \omega))\|_{k, \epsilon} dP(\omega) < \infty$$

for any fixed $0 < \rho, a < \infty$ and $\epsilon \in (0, 1]$.

Define the random field $Z : \mathbf{R}^+ \times H \times \Omega \rightarrow H$ by

$$Z(t, x, \omega) := U(t, x + Y(\omega), \omega) - Y(\theta(t, \omega))$$

for $t \geq 0, x \in H, \omega \in \Omega$. Then (Z, θ) is a $C^{k,\epsilon}$ perfect cocycle. Furthermore, there is a sure event $\Omega_5 \in \mathcal{F}$ with the following properties:

(i) $\theta(t, \cdot)(\Omega_5) = \Omega_5$ for all $t \in \mathbf{R}$,

(ii) For every $\omega \in \Omega_5$ and any $x \in H$, the statement

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)| < 0 \quad (3.23)$$

implies

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |Z(t, x, \omega)| = \limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)|. \quad (3.24)$$

Proof.

Note that, by definition, Z is a “centering” of the cocycle U with respect to the stationary trajectory $\{Y(\theta(t, \cdot)) : t \geq 0\}$ in the sense that $Z(t, 0, \omega) = 0$ for all $(t, \omega) \in \mathbf{R}^+ \times \Omega$. Furthermore, (Z, θ) is a $C^{k,\epsilon}$ perfect cocycle. To see this let $t_1, t_2 \geq 0, \omega \in \Omega, x \in H$. Then by the perfect cocycle property for U , it follows that

$$\begin{aligned} Z(t_2, Z(t_1, x, \omega), \theta(t_1, \omega)) &= U(t_2, Z(t_1, x, \omega) + Y(\theta(t_1, \omega)), \theta(t_1, \omega)) - Y(\theta(t_2, \theta(t_1, \omega))) \\ &= U(t_2, U(t_1, x + Y(\omega), \omega), \theta(t_1, \omega)) - Y(\theta(t_2 + t_1, \omega)) \\ &= Z(t_1 + t_2, x, \omega). \end{aligned}$$

Using the integrability condition of the lemma, the proofs of assertions (i) and (ii) follow in the same manner as for the corresponding assertions in Lemma 3.4 ([M-S.3]). $\square$

Proof of Theorem 2.1.

The proof of the theorem consists in two major undertakings:

- (a) Using Ruelle’s discrete-time analysis [Ru.2] to show that the assertions of Theorem 2.1 hold for the discretized cocycle, perfectly in ω .
- (b) Extending the discrete-time results to continuous time via perfection techniques that are essentially based on the ergodic theorem and Kingman’s subadditive ergodic theorem.

Recall the auxiliary cocycle (Z, θ) defined in Lemma 3.4. Consider the random family of maps $F_\omega : \bar{B}(0, 1) \rightarrow H$, $\omega \in \Omega$, given by $F_\omega(x) := Z(1, x, \omega)$, $x \in H$, and the time-one shift $\tau := \theta(1, \cdot) : \Omega \rightarrow \Omega$. Adopting Ruelle's notation ([Ru.2], p. 272), we set $F_\omega^n := F_{\tau^{n-1}(\omega)} \circ \dots \circ F_{\tau(\omega)} \circ F_\omega$. Therefore, $F_\omega^n = Z(n, \cdot, \omega)$ for each $n \geq 1$, because (Z, θ) is a cocycle. By Lemma 3.4, each map F_ω is $C^{k, \epsilon}$ ($\epsilon \in (0, 1]$) on $\bar{B}(0, 1)$ and by the definition of Z , it follows that $(DF_\omega)(0) = DU(1, Y(\omega), \omega)$. By the integrability hypothesis of the theorem, it is clear that $\log^+ \|DU(1, Y(\cdot), \cdot)\|_{L(H)}$ is integrable. Moreover, in view of the integrability hypothesis on (U, θ) , it follows that the linearized continuous-time cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$ and the discrete-time cocycle $((DF_\omega^n)(0), \theta(n, \omega))$ share the same Lyapunov spectrum, viz.:

$$\{-\infty < \dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}.$$

(Cf. [Mo.1]). Assume that λ_{i_0} is finite, that is $\lambda_{i_0} \in (-\infty, 0)$. Therefore, under hypotheses (I) of Theorem 5.1 in ([Ru.2], p. 272), there is a sure event $\Omega_1^* \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_1^*) = \Omega_1^*$ for all $t \in \mathbf{R}$, $\bar{\mathcal{F}}$ -measurable positive random variables $\rho_1, \beta_1 : \Omega_1^* \rightarrow (0, 1)$, and a random family of $C^{k, \epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) stable submanifolds $\tilde{\mathcal{S}}_d(\omega)$ of $\bar{B}(0, \rho_1(\omega))$ satisfying the following properties for each $\omega \in \Omega_1^*$:

$$\tilde{\mathcal{S}}_d(\omega) = \{x \in \bar{B}(0, \rho_1(\omega)) : |Z(n, x, \omega)| \leq \beta_1(\omega)e^{(\lambda_{i_0} + \epsilon_1)n} \text{ for all integers } n \geq 0\}. \quad (3.25)$$

When $\lambda_{i_0} = -\infty$, the stable manifold is defined by

$$\tilde{\mathcal{S}}_d(\omega) = \{x \in \bar{B}(0, \rho_1(\omega)) : |Z(n, x, \omega)| \leq \beta_1(\omega)e^{\lambda n} \text{ for all integers } n \geq 0\}, \quad (3.25')$$

where $\lambda \in (-\infty, 0)$ is arbitrary. The stable subspace $\mathcal{S}(\omega)$ of the linearized cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$ is tangent to the stable manifold $\tilde{\mathcal{S}}_d(\omega)$ at 0; viz. $T_0\tilde{\mathcal{S}}_d(\omega) = \mathcal{S}(\omega)$. In particular, $\text{codim } \tilde{\mathcal{S}}_d(\omega)$ is finite and non-random. Again by Theorem 5.1 of [Ru.2]), we have the following estimate on the Lyapunov exponent of the Lipschitz constant of $(Z(n, \cdot), \theta(n, \cdot))$ over its stable manifold:

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \left[\sup_{\substack{x_1, x_2 \in \tilde{\mathcal{S}}_d(\omega) \\ x_1 \neq x_2}} \frac{|Z(n, x_1, \omega) - Z(n, x_2, \omega)|}{|x_1 - x_2|} \right] \leq \lambda_{i_0}. \quad (3.26)$$

The statements in the above paragraph hold for all ω in the $\theta(t, \cdot)$ -invariant sure event Ω_1^* . In order to construct such an event, we will use perfection arguments and the proof of Theorem 5.1 ([Ru.2], p. 272). Assume first that $k = 1$ and $\epsilon > 0$. Using the notation of [Ru.2], denote $T^t(\omega) := DZ(t, 0, \omega)$, $f(\omega) := \theta(1, \omega)$, $T_n(\omega) := DZ(1, 0, \theta((n-1), \omega))$, for all $\omega \in \Omega$, any positive real t and any integer $n \geq 1$. It is possible to replace (5.3) in [Ru.2], p. 274) by its continuous-time *perfect* analogue

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log^+ \|Z(1, \cdot, \theta(t, \omega))\|_{1, \epsilon} = 0. \quad (3.27)$$

This follows from the integrability hypothesis of the theorem and the perfect ergodic theorem (Lemma 3.1 (ii)). More specifically, (3.27) holds for all ω in a sure event $\Omega_1^* \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_1^*) = \Omega_1^*$ for all $t \in \mathbf{R}$. Assume $\lambda_{i_0} > -\infty$. Adopting the terminology of Theorem 1.1 ([Ru.2], p. 248), take $S = i_0 - 1$, and $\mu^{(S+1)} = \lambda_{i_0}$. In case $\lambda_{i_0} = -\infty$, take $\mu^{(S+1)}$ to be any fixed number in $(-\infty, 0)$. The integrability hypothesis on U coupled with Lemma 3.3 (where $j_0 = i_0 - 1$) imply the existence of a sure event $\Omega_2^* \in \mathcal{F}$ such that $\Omega_2^* \subseteq \Omega_1^*$, $\theta(t, \cdot)(\Omega_2^*) = \Omega_2^*$ for all $t \in \mathbf{R}$, and the sequence $T_n(\omega), V_n(\omega) := E_{i_0}(\tau^n(\omega))$, $n \geq 1$, satisfies Conditions (S) of [Ru.2], p. 256) for every $\omega \in \Omega_2^*$. Pick and fix any $\omega \in \Omega_2^*$. As in the proof of Theorem 5.1 ([Ru.2], pp. 274-278), the “perturbation theorem” (Theorem 4.1, [Ru.2], pp. 262-263) holds for the sequence $T_n(\omega), n \geq 1$. Thus the assertions in the previous paragraph are valid for $k = 1$ and any $\epsilon \in (0, 1]$. When $k > 1$ and $\epsilon \in (0, 1]$, we first apply the previous analysis to the perfect cocycle

$$\left(\check{Z}(t, x, x_1, \omega) := (Z(t, x, \omega), DZ(t, x, \omega)x_1, \theta(t, \omega)) \right), \quad x, x_1 \in H, t \geq 0,$$

on $H \oplus H$. Secondly, we use the inductive argument of ([Ru.2], pp. 278-279) to show that the $\tilde{\mathcal{S}}_d(\omega)$ are $C^{k, \epsilon}$ manifolds ($k > 1, \epsilon \in (0, 1]$) *perfectly in* ω .

To establish assertion (a) of the theorem, let $\tilde{\mathcal{S}}(\omega), \omega \in \Omega_1^*$, be the set defined therein. Then the definition of Z and property (3.25) of $\tilde{\mathcal{S}}_d(\omega)$ imply that

$$\tilde{\mathcal{S}}(\omega) = \tilde{\mathcal{S}}_d(\omega) + Y(\omega) \quad (3.28)$$

for all $\omega \in \Omega_1^*$. Thus $\tilde{\mathcal{S}}(\omega)$ is a $C^{k,\epsilon}$ manifold ($k > 1, \epsilon \in (0, 1]$), with tangent space $T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = T_0\tilde{\mathcal{S}}_d(\omega) = \mathcal{S}(\omega)$. In particular, $\text{codim } \tilde{\mathcal{S}}(\omega) = \text{codim } \mathcal{S}(\omega), \omega \in \Omega_1^*$, is finite and non-random.

To complete the proof of the inequality (2.1) in part (a) of the theorem, use (3.26) to get

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)| \leq \lambda_{i_0}$$

perfectly in ω for all $x \in \tilde{\mathcal{S}}_d(\omega)$. In view of Lemma 3.4, we may extend the above estimate to cover its continuous-time counterpart. Hence we obtain a sure event $\Omega_3^* \subseteq \Omega_2^*, \Omega_3^* \in \mathcal{F}$, such that $\theta(t, \cdot)(\Omega_3^*) = \Omega_3^*$ for all $t \in \mathbf{R}$, and

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |Z(t, x, \omega)| \leq \lambda_{i_0} \quad (3.29)$$

for all $\omega \in \Omega_3^*$ and all $x \in \tilde{\mathcal{S}}_d(\omega)$. The above inequality together with definition of Z imply the estimate (2.1) of the theorem.

Next, we establish assertion (b) of the theorem. To do so, let $\omega \in \Omega_1^*$ and $x \in \tilde{\mathcal{S}}_d(\omega)$. Then by (3.26), it follows that there is a positive integer $N_0 := N_0(\omega)$, independent of $x \in \tilde{\mathcal{S}}_d(\omega)$, such that $Z(n, x, \omega) \in \bar{B}(0, 1)$ for all $n \geq N_0$. Now Lemma 3.1(ii) gives a $\theta(t, \cdot)$ -invariant sure event Ω_3 such that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log^+ \sup_{\substack{0 \leq u \leq 1, \\ (v^*, \eta^*) \in \bar{B}(0, 1)}} \|DZ(u, (v^*, \eta^*), \theta(t, \omega))\|_{L(H)} = 0 \quad (3.29')$$

for all $\omega \in \Omega_3$. Define the sure event $\Omega_4^* := \Omega_3^* \cap \Omega_3 \in \mathcal{F}$. Clearly, $\theta(t, \cdot)(\Omega_4^*) = \Omega_4^*$ for all $t \in \mathbf{R}$.

By the definition of Z and the Mean Value Theorem, we obtain the following inequalities

$$\begin{aligned} & \sup_{n \leq t \leq n+1} \frac{1}{t} \log \left[\sup_{\substack{x_1 \neq x_2, \\ (v_1, \eta_1), x_2 \in \tilde{\mathcal{S}}(\omega)}} \frac{|U(t, x_1, \omega) - U(t, x_2, \omega)|}{|x_1 - x_2|} \right] \\ &= \sup_{n \leq t \leq n+1} \frac{1}{t} \log \left[\sup_{\substack{x_1 \neq x_2, \\ x_1, x_2 \in \tilde{\mathcal{S}}_d(\omega)}} \frac{|Z(t, x_1, \omega) - Z(t, x_2, \omega)|}{|x_1 - x_2|} \right] \\ &\leq \frac{1}{n} \log^+ \sup_{\substack{0 \leq u \leq 1, \\ (v^*, \eta^*) \in \bar{B}(0, 1)}} \|DZ(u, (v^*, \eta^*), \theta(n, \omega))\|_{L(H)} \\ &\quad + \frac{n}{(n+1)} \frac{1}{n} \log \left[\sup_{\substack{x_1 \neq x_2, \\ x_1, x_2 \in \tilde{\mathcal{S}}_d(\omega)}} \frac{|Z(n, x_1, \omega) - Z(n, x_2, \omega)|}{|x_1 - x_2|} \right] \end{aligned}$$

for all $\omega \in \Omega_4^*$, and all sufficiently large $n \geq N_0(\omega)$. Now take $\limsup_{n \rightarrow \infty}$ on both sides of the above inequality, and use (3.26), (3.29') in order to complete the proof assertion (b) of the theorem.

The cocycle invariance (2.3) in part (c) of the theorem follows immediately from the Oseledec-Ruelle theorem (Theorem 1.1) applied to the perfect linearized cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$. Indeed, one gets a sure $\theta(t, \cdot)$ -invariant event $\in \mathcal{F}$ (also denoted by Ω_1^*), such that $DU(t, Y(\omega), \omega)(\mathcal{S}(\omega)) \subseteq \mathcal{S}(\theta(t, \omega))$ for all $t \geq 0$ and all $\omega \in \Omega_1^*$.

The proof of the asymptotic invariance property (2.2) of the non-linear cocycle requires some work. To achieve this, we will extend the arguments underlying the proofs of Theorems 5.1 and 4.1 in [Ru.2], pp. 262-279, to a continuous time setting. The crucial step towards this goal is to show that the two random variables ρ_1, β_1 in (3.25) may be redefined on a sure event (also denoted by) Ω_1^* such that $\theta(t, \cdot)(\Omega_1^*) = \Omega_1^*$ for all $t \in \mathbf{R}$, and

$$\rho_1(\theta(t, \omega)) \geq \rho_1(\omega)e^{(\lambda_{i_0} + \epsilon_1)t}, \quad \beta_1(\theta(t, \omega)) \geq \beta_1(\omega)e^{(\lambda_{i_0} + \epsilon_1)t} \quad (3.30)$$

for every $\omega \in \Omega_1^*$ and all $t \geq 0$. For the given choice of ϵ_1 , fix $0 < \epsilon_3 < -\epsilon(\lambda_{i_0} + \epsilon_1)/4$, where $\epsilon \in (0, 1]$ denotes the Hölder exponent of U . The above inequalities hold in the *discrete* case (when $t = n$, a positive integer) because of Theorem 5.1 (c) ([Ru.2], p. 274). To prove them for any continuous time t , we will modify the definitions of ρ_1, β_1 in the proofs of Theorems 5.1 and 4.1 in [Ru.2]. In the notation of the proof of Theorem 5.1 ([Ru.2], p. 274), we replace the random variable G in (5.4) ([Ru.2], p. 274) by the larger one

$$\tilde{G}(\omega) := \sup_{t \geq 0} \|Z(1, \cdot, \theta(t, \omega))\|_{1, \epsilon} e^{(-t\epsilon_3 - \lambda\epsilon)}. \quad (3.31)$$

Clearly, $\tilde{G}(\omega)$ is finite perfectly in ω , because of (3.27) and Lemma 3.2. Following ([Ru.2], pp. 266, 274), the random variables ρ_1, β_1 may be chosen according to the relations

$$\beta_1 := \left[\frac{\delta_1 \wedge \left(\frac{1}{\sqrt{2A}} \right)}{2\tilde{G}} \right]^{\frac{1}{\epsilon}} \wedge 1 \quad (3.32)$$

$$\rho_1 := \frac{\beta_1}{B_{\epsilon_3}} \quad (3.33)$$

where A, δ_1 and B_{ϵ_3} are random positive constants that are defined via continuous-time analogues of the relations (4.26), (4.18)-(4.21), (4.24), (4.25) in [Ru.2], pp. 265-267, with η replaced by ϵ_3 . In particular, the “ancestry” of A, δ_1 and B_{ϵ_3} in Ruelle’s argument may be traced back to the constants $D_{\epsilon_3}, K_{\epsilon_3}$ which appear in Lemmas 3.3 and 3.2 of this article. Hence (3.30) will follow if we can show that, for sufficiently small $\epsilon_3 > 0$, the following inequalities

$$\left. \begin{aligned} K_{\epsilon_3}(\theta(l, \omega)) &\leq K_{\epsilon_3}(\omega) + \frac{\epsilon_3 l}{2} \\ D_{\epsilon_3}(\theta(l, \omega)) &\leq e^{\frac{\epsilon_3 l}{2}} D_{\epsilon_3}(\omega) \\ \tilde{G}(\theta(l, \omega)) &\leq e^{\epsilon_3 l} \tilde{G}(\omega) \end{aligned} \right\} \quad (3.34)$$

hold perfectly in ω for all real $l \geq 0$. The first and second inequalities in (3.34) follow from Lemmas 3.2(ii) and 3.3, respectively. The third inequality is an immediate consequence of the definition of $\tilde{G}$ in (3.31). The proof of (3.30) is now complete in view of (3.32), (3.33) and (3.34).

The inequalities in (3.30) will allow us to establish the asymptotic invariance property (2.2) in (c) of the theorem. By the perfect inequality in (b), there is a sure event $\Omega_5^* \subseteq \Omega_4^*$ such that $\theta(t, \cdot)(\Omega_5^*) = \Omega_5^*$ for all $t \in \mathbf{R}$, and for any $0 < \epsilon' < \epsilon_1$ and any $\omega \in \Omega_5^*$, there exists $\beta^{\epsilon'}(\omega) > 0$ (independent of x) so that

$$|U(t, x, \omega) - Y(\theta(t, \omega))| \leq \beta^{\epsilon'}(\omega) e^{(\lambda_{i_0} + \epsilon')t} \quad (3.35)$$

for all $x \in \tilde{\mathcal{S}}(\omega)$, $t \geq 0$. Let t be any positive real, n a non-negative integer, $\omega \in \Omega_5^*$ and $x \in \tilde{\mathcal{S}}(\omega)$.

Using the cocycle property and (3.35), we obtain

$$\begin{aligned} |U(n, U(t, x, \omega), \theta(t, \omega)) - Y(\theta(n, \theta(t, \omega)))| &= |U(n+t, x, \omega) - Y(\theta(n+t, \omega))| \\ &\leq \beta^{\epsilon'}(\omega) e^{(\lambda_{i_0} + \epsilon')(n+t)} \\ &\leq \beta^{\epsilon'}(\omega) e^{(\lambda_{i_0} + \epsilon')t} e^{(\lambda_{i_0} + \epsilon_1)n}. \end{aligned} \quad (3.36)$$

Using (3.30), (3.35), (3.36) and the definition of $\tilde{\mathcal{S}}(\theta(t, \omega))$, we see that for each $\omega \in \Omega_5^*$, there exists $\tau_1(\omega) > 0$ such that $U(t, x, \omega) \in \tilde{\mathcal{S}}(\theta(t, \omega))$ for all $t \geq \tau_1(\omega)$. Hence, for all $\omega \in \Omega_5^*$,

$$U(t, \cdot, \omega)(\tilde{\mathcal{S}}(\omega)) \subseteq \tilde{\mathcal{S}}(\theta(t, \omega)), \quad t \geq \tau_1(\omega)$$

and the proof of assertion (c) is complete.

Our next objective is to establish the existence of the *perfect* family of local unstable manifolds $\tilde{\mathcal{U}}(\omega)$ in assertion (d) of the theorem. To this end, we define the random field $\hat{Z} : \mathbf{R}^+ \times H \times \Omega \rightarrow H$ by

$$\hat{Z}(t, x, \omega) := U(t, x + Y(\theta(-t, \omega)), \theta(-t, \omega)) - Y(\omega) \quad (3.37)$$

for all $t \geq 0$, $x \in H$, $\omega \in \Omega$. Note that $\hat{Z}(t, \cdot, \omega) = Z(t, \cdot, \theta(-t, \omega))$, $t \geq 0$, $\omega \in \Omega$; and $\hat{Z}$ is $(\mathcal{B}(\mathbf{R}^+) \otimes \mathcal{B}(H) \otimes \mathcal{F}, \mathcal{B}(H))$ -measurable. Since Y is a stationary point for (U, θ) , we may replace ω by $\theta(-t, \omega)$ in (1.1). Thus $\hat{Z}(t, 0, \omega) = 0$ for all $t \geq 0$, $\omega \in \Omega$. We contend that $([D\hat{Z}(t, 0, \omega)]^*, \theta(-t, \omega), t \geq 0)$ is a perfect linear cocycle in $L(H)$. To see this, we first observe that $(DU(t, Y(\omega), \omega), \theta(t, \omega))$ is an $L(H)$ -valued perfect cocycle:

$$DU(t_1 + t_2, Y(\omega), \omega) = DU(t_1, Y(\theta(t_2, \omega)), \theta(t_2, \omega)) \circ DU(t_2, Y(\omega), \omega)$$

for all $\omega \in \Omega, t_1, t_2 \geq 0$. Secondly, we replace ω by $\theta(-t_1 - t_2, \omega)$ and take adjoints in the above identity to obtain

$$\begin{aligned} & [DU(t_1 + t_2, Y(\theta(-t_1 - t_2, \omega)), \theta(-t_1 - t_2, \omega))]^* \\ &= [DU(t_2, Y(\theta(-t_1 - t_2, \omega)), \theta(-t_1 - t_2, \omega))]^* \circ [DU(t_1, Y(\theta(-t_1, \omega)), \theta(-t_1, \omega))]^* \end{aligned}$$

for all $\omega \in \Omega, t_1, t_2 \geq 0$. Therefore,

$$[D\hat{Z}(t_1 + t_2, 0, \omega)]^* = [D\hat{Z}(t_2, 0, \theta(-t_1, \omega))]^* \circ [D\hat{Z}(t_1, 0, \omega)]^*$$

for all $\omega \in \Omega, t_1, t_2 \geq 0$; and our contention is proved.

We will now apply the Oseledec-Ruelle theorem to the perfect cocycle $([D\hat{Z}(t, 0, \omega)]^*, \theta(-t, \omega), t \geq 0)$. To do this, it is sufficient to check the integrability condition

$$\int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \|[D\hat{Z}(t_2, 0, \theta(-t_1, \omega))]^*\|_{L(H)} dP(\omega) < \infty \quad (3.38)$$

for any fixed $a \in (0, \infty)$. The above integrability relation follows from the integrability hypothesis of Theorem 2.1 and the P -preserving property of $\theta(t, \cdot)$:

$$\begin{aligned}
& \int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \| [D\hat{Z}(t_2, 0, \theta(-t_1, \omega))]^* \|_{L(H)} dP(\omega) \\
&= \int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \| DU(t_2, Y(\theta(-t_2 - t_1, \omega)), \theta(-t_2 - t_1, \omega)) \|_{L(H)} dP(\omega) \\
&\leq \int_{\Omega} \log^+ \sup_{0 \leq t_1 \leq 2a, 0 \leq t_2 \leq a} \| DU(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega)) \|_{L(H)} dP(\omega) \\
&\leq \int_{\Omega} \log^+ \sup_{0 \leq t_1 \leq a, 0 \leq t_2 \leq a} \| DU(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega)) \|_{L(H)} dP(\omega) \\
&+ \int_{\Omega} \log^+ \sup_{a \leq t_1 \leq 2a, 0 \leq t_2 \leq a} \| DU(t_2, Y(\theta(t_1 - a, \omega)), \theta(t_1 - a, \omega)) \|_{L(H)} dP(\omega) \\
&= 2 \int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \| DU(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega)) \|_{L(H)} dP(\omega) < \infty.
\end{aligned}$$

By (3.38) and the Oseledec-Ruelle theorem, we conclude that the linear cocycle

$([D\hat{Z}(t, 0, \omega)]^*, \theta(-t, \omega), t \geq 0)$ has a fixed discrete Lyapunov spectrum. Furthermore, this spectrum (with multiplicities) coincides with that of the cocycle $(DU(t, Y(\omega), \omega), \theta(t, \omega))$, viz. $\{\dots \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$ where $\lambda_i \neq 0$ for all $i \geq 1$, by hyperbolicity. See [Ru.2], Section 3.5, p. 261.

The next step in our construction of the *perfect* random family of local unstable manifolds $\tilde{\mathcal{U}}(\omega)$ is the following estimate:

$$\int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq 1} \| \hat{Z}(t_2, \cdot, \theta(-t_1, \omega)) \|_{k, \epsilon} dP(\omega) < \infty.$$

By the same argument as in the previous paragraph, the above estimate is a consequence of the P -preserving property of $\theta(t, \cdot)$, $t \in \mathbf{R}$, and the integrability hypothesis of the theorem. Define λ_{i_0-1} as in the statement of Theorem 2.1, and fix any $\epsilon_2 \in (0, \lambda_{i_0-1})$. In view of the above integrability property, it follows from Lemma 3.3 that the sequence $\tilde{T}_n(\omega) := [D\hat{Z}(1, 0, \theta(-n, \omega))]^*, \theta(-n, \omega)$, $n \geq 0$, satisfies Condition (S) of [Ru.2] perfectly in ω . Hence the sequence $\tilde{T}_n(\omega)$, $n \geq 1$, satisfies Corollary 3.4 ([Ru.2], p. 260) perfectly in ω , because of Proposition 3.3 in [Ru.2]. At this point, we may modify the arguments in the proof of Ruelle's Theorem 6.1 ([Ru.2], p. 280) using an approach analogous to the one used in constructing the stable manifolds in this proof. Therefore, one gets a $\theta(-t, \cdot)$ -invariant sure event $\hat{\Omega}_1^* \in \mathcal{F}$ and $\bar{\mathcal{F}}$ -measurable random variables $\rho_2, \beta_2 : \hat{\Omega}_1^* \rightarrow (0, 1)$

satisfying the following properties. If $\lambda_{i_0-1} < \infty$, define $\tilde{\mathcal{U}}_d(\omega)$ to be the set of all $x_0 \in \bar{B}(0, \rho_2(\omega))$ with the property that there is a discrete “history” process $u(-n, \cdot) : \Omega \rightarrow H, n \geq 0$, such that $u(0, \omega) = x_0$, $\hat{Z}(1, u(-(n+1), \omega), \theta(-n, \omega)) = u(-n, \omega)$ and $|u(-n, \omega)| \leq \beta_2(\omega)e^{-n(\lambda_{i_0-1}-\epsilon_2)}$ for all $n \geq 0$. If $\lambda_{i_0-1} = \infty$, define $\tilde{\mathcal{U}}_d(\omega)$ to be the set of all $x_0 \in H$ with the property that there is a discrete history process $u(-n, \cdot) : \Omega \rightarrow H, n \geq 0$, such that $u(0, \omega) = x_0$, and $|u(-n, \omega)| \leq \beta_2(\omega)e^{-\lambda n}$ for all $n \geq 0$ and arbitrary $\lambda > 0$. It follows from ([Ru.2], p. 281) that the discrete history process $u(-n, \cdot)$ is uniquely determined by x_0 . Moreover, each $\tilde{\mathcal{U}}_d(\omega), \omega \in \hat{\Omega}_1^*$, is a $C^{k, \epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) finite-dimensional submanifold of $\bar{B}(0, \rho_2(\omega))$ with tangent space $\mathcal{U}(\omega)$ at 0, and $\dim \tilde{\mathcal{U}}_d(\omega)$ is fixed independently of ω and ϵ_2 . Furthermore,

$$\rho_2(\theta(-t, \omega)) \geq \rho_2(\omega)e^{-(\lambda_{i_0-1}-\epsilon_2)t}, \quad \beta_2(\theta(-t, \omega)) \geq \beta_2(\omega)e^{-(\lambda_{i_0-1}-\epsilon_2)t}. \quad (3.39)$$

perfectly in ω for all $t \geq 0$. We claim that the set $\tilde{\mathcal{U}}(\omega)$ defined in (d) of Theorem 2.1 coincides with $\tilde{\mathcal{U}}_d(\omega) + Y(\omega)$ for each $\omega \in \hat{\Omega}_1^*$. We first show that $\tilde{\mathcal{U}}_d(\omega) + Y(\omega) \subseteq \tilde{\mathcal{U}}(\omega)$. Let $x_0 \in \tilde{\mathcal{U}}_d(\omega)$ and u be as above. Set

$$y_0(-n, \omega) := u(-n, \omega) + Y(\theta(-n, \omega)), \quad n \geq 0. \quad (3.40)$$

It is easy to check that y_0 is a discrete history process satisfying the first and second assertions in (d) of the theorem. Hence $x_0 + Y(\omega) \in \tilde{\mathcal{U}}(\omega)$. Similarly, $\tilde{\mathcal{U}}(\omega) \subseteq \tilde{\mathcal{U}}_d(\omega) + Y(\omega)$ for all $\omega \in \hat{\Omega}_1^*$. Hence $\tilde{\mathcal{U}}(\omega) = \tilde{\mathcal{U}}_d(\omega) + Y(\omega)$ for all $\omega \in \hat{\Omega}_1^*$. This immediately implies that $\tilde{\mathcal{U}}(\omega)$ is a $C^{k, \epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) finite-dimensional submanifold of $\bar{B}(Y(\omega), \rho_2(\omega))$, and

$$T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) = T_0\tilde{\mathcal{U}}_d(\omega) = \mathcal{U}(\omega)$$

for all $\omega \in \hat{\Omega}_1^*$.

We will next address the issue of the existence of the continuous-time history process satisfying the third assertion in part (d) of the theorem. Suppose $x \in \tilde{\mathcal{U}}(\omega)$. From what we proved in the previous paragraph, it follows that there is an $x_0 \in \mathcal{U}_d(\omega)$ such that $x = x_0 + Y(\omega)$. The discrete process y_0 given by (3.40) may be extended to a continuous-time history process $y(\cdot, \omega) : (-\infty, 0] \rightarrow H$ such that $y(0, \omega) = x$, and $y(\cdot, \omega)$ satisfies the third assertion in (d). This is

achieved by interpolation within the periods $[-(n+1), -n]$, $n \geq 0$, using the cocycle property of U : Indeed, let $s \in (-(n+1), -n)$. Then there is an $\alpha \in (0, 1)$, such that $s = \alpha - (n+1)$. Define

$$y(s, \omega) := U(s + n + 1, y_0(-(n+1), \omega), \theta(-(n+1), \omega)).$$

Obviously, $y(0, \omega) = x_0 + Y(\omega) = x$. Let $s \in (-(n+1), -n)$ and suppose $0 < t \leq -s$. Pick a positive integer $m < n$ such that $s + t \in [-(m+1), -m]$. The above definition of y , together with the perfect cocycle property for U , easily imply that

$$y(t + s, \omega) = U(t, y(s, \omega), \theta(s, \omega)). \quad (3.41)$$

In particular, $U(t, y(-t, \omega), \theta(-t, \omega)) = x$ for all $t \geq 0$. This follows from (3.41) when s is replaced by $-t$. Furthermore, for each $x \in \tilde{\mathcal{U}}(\omega)$, the above continuous-time history process is uniquely determined because its discrete-time counterpart is unique.

We will now prove the following estimate

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |y(-t, \omega) - Y(\theta(-t, \omega))| \leq -\lambda_{i_0-1} \quad (3.42)$$

perfectly in ω . We start with its discrete-time counterpart

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |y(-n, \omega) - Y(\theta(-n, \omega))| \leq -\lambda_{i_0-1} \quad (3.43)$$

which holds perfectly in ω , because of Theorem 6.1 (b) in [Ru.2]. Let $t \in (n, n+1)$. Then there exists $\gamma \in (0, 1)$ such that $-t = \gamma - (n+1)$. Thus by the definition of y and the Mean Value Theorem, it follows that

$$\begin{aligned} & |y(-t, \omega) - Y(\theta(-t, \omega))| \\ &= |U(\gamma, y(-(n+1), \omega), \theta(-(n+1), \omega)) - U(\gamma, Y(\theta(-(n+1), \omega), \theta(-(n+1), \omega)))| \\ &\leq \sup_{\substack{(v^*, \eta^*) \in \tilde{B}(0, 1), \\ \gamma \in (0, 1)}} \|DU(\gamma, (v^*, \eta^*) + Y(\theta(-(n+1), \omega)), \theta(-(n+1), \omega))\|_{L(H)} \\ &\quad \times |y(-(n+1), \omega) - Y(\theta(-(n+1), \omega))| \end{aligned}$$

perfectly in ω . Letting $t \rightarrow \infty$, we get

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log |y(-t, \omega) - Y(\theta(-t, \omega))| \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log^+ \sup_{\substack{(v^*, \eta^*) \in \bar{B}(0, 1), \\ \gamma \in (0, 1)}} \|DU(\gamma, (v^*, \eta^*) + Y(\theta(-(n+1), \omega)), \theta(-(n+1), \omega))\|_{L(H)} \\ & \quad + \limsup_{n \rightarrow \infty} \frac{1}{n} \log |y(-(n+1), \omega) - Y(\theta(-(n+1), \omega))|. \end{aligned}$$

By the integrability condition of the theorem and the perfect ergodic theorem (Lemma 3.1 (ii)), the first term on the right hand side of the above inequality is zero, perfectly in $\omega \in \Omega$. Since $y(0) \in \tilde{\mathcal{U}}(\omega)$, the second term is less than or equal to $-\lambda_{i_0-1}$. This completes the proof of assertion (d) of the theorem.

We will omit the proof of assertion (e), since it is very similar to that of (3.42).

Our next objective is to prove assertion (f) of the theorem. Note first that the perfect invariance

$$DU(t, \cdot, \theta(-t, \omega))(\mathcal{U}(\theta(-t, \omega))) = \mathcal{U}(\omega), \quad t \geq 0,$$

follows from the cocycle property for the linearized semiflow and Theorem (1.2); cf. [Mo.1], Corollary 2 (v) of Theorem 4. Since $\dim \mathcal{U}(\omega)$ is fixed and finite perfectly in ω , the restriction

$$DU(t, \cdot, \theta(-t, \omega))|_{\mathcal{U}(\theta(-t, \omega))} : \mathcal{U}(\theta(-t, \omega)) \rightarrow \mathcal{U}(\omega), \quad t \geq 0,$$

is a linear homeomorphism onto. It remains to check the following asymptotic invariance property in (f):

$$\tilde{\mathcal{U}}(\omega) \subseteq U(t, \cdot, \theta(-t, \omega))(\tilde{\mathcal{U}}(\theta(-t, \omega))), \quad t \geq \tau_2(\omega), \quad (3.44)$$

perfectly in ω for some $\tau_2(\omega) > 0$. Suppose $x \in \tilde{\mathcal{U}}(\omega)$. Then by assertions (d), (e) of the theorem and inequalities (3.39), there exist a (unique) history process $y(-t, \omega), t \geq 0$, and a random time $\tau_2(\omega) > 0$ satisfying the following: $y(0, \omega) = x$, $y(-t, \omega) \in \bar{B}(Y(\theta(-t, \omega)), \rho_2(\theta(-t, \omega)))$ for all $t \geq \tau_2(\omega)$, and

$$y(t' - t, \omega) = U(t', y(-t, \omega), \theta(-t, \omega)), \quad 0 < t' \leq t, \quad (3.45)$$

perfectly in ω . Pick any $t_1 \geq \tau_2(\omega)$. Then $x = U(t_1, y(-t_1, \omega), \theta(-t_1, \omega))$, because of (3.45) (for $t = t' = t_1$). Now $y(-t_1, \omega) \in \tilde{\mathcal{U}}(\theta(-t_1, \omega))$. To prove this, we define the process $y_1(-t, \omega) := y(-t - t_1, \omega)$, $t \geq 0$. Hence $y_1(\cdot, \omega)$ is a history process and

$$y_1(0, \omega) = y(-t_1, \omega) \in \bar{B}(Y(\theta(-t_1, \omega)), \rho_2(\theta(-t_1, \omega))).$$

Therefore $y(-t_1, \omega) \in \tilde{\mathcal{U}}(\theta(-t_1, \omega))$. This implies (3.44) because $t_1 \geq \tau_2(\omega)$ is arbitrary.

To prove the transversality property in (g), note the following perfect identities:

$$T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) = \mathcal{U}(\omega), \quad T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = \mathcal{S}(\omega), \quad H = \mathcal{U}(\omega) \oplus \mathcal{S}(\omega).$$

All the assertions (a)-(g) of the theorem will hold perfectly in ω if we take $\Omega^* := \Omega_1^* \cap \hat{\Omega}_1^*$.

To deal with the case when U is a C^∞ cocycle, we adapt the proof in [Ru.2], section (5.3) (p. 297). Thus we obtain a $\theta(t, \cdot)$ -invariant sure event in $\mathcal{F}$ (also denoted by Ω^*) such that $\tilde{\mathcal{S}}(\omega)$ and $\tilde{\mathcal{U}}(\omega)$ are C^∞ for all $\omega \in \Omega^*$. This completes the proof of Theorem 2.1. $\square$

4. The local stable manifold theorem for see's and spde's.

In this section, we discuss several classes of semilinear stochastic evolutions equations and spde's. The objective is to establish sufficient conditions for a local stable manifold theorem for each class.

(a) *Stochastic semilinear evolution equations: Additive noise.*

Let K, H be two separable real Hilbert spaces. Let A be a self-adjoint operator on H such that $A \geq cI_H$, where c is a real constant and I_H is the identity operator on H . Assume that A admits a discrete non-vanishing spectrum $\{\mu_n, n \geq 1\}$ which is bounded below. Let $\{e_n, n \geq 1\}$ denote a basis for H consisting of eigen vectors of A , viz. $Ae_n = \mu_n e_n$, $n \geq 1$. Assume further that A^{-1} is trace-class. Suppose $B_0 \in L_2(K, H)$. Let $W(t), t \in \mathbf{R}$, be cylindrical Brownian motion on the canonical filtered Wiener space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, P)$ and with a separable covariance Hilbert space K ([M.Z.Z.1], section 2). Let $T_t = e^{-At}$ stand for the strongly continuous semigroup generated by $-A$.

Denote by μ_m the largest negative eigenvalue of A and by μ_{m+1} its smallest positive eigenvalue. Thus there is an orthogonal $\{T_t\}_{t \geq 0}$ -invariant splitting of H using the negative eigenvalues $\{\mu_1, \mu_2, \dots, \mu_m\}$ and the positive eigenvalues $\{\mu_n : n \geq m+1\}$ of A :

$$H = H^+ \oplus H^-$$

where H^+ is a closed linear subspace of H and H^- is a finite-dimensional subspace. Denote by $p^+ : H \rightarrow H^+$ and $p^- : H \rightarrow H^-$ the corresponding projections onto H^+ and H^- respectively. Since H^- is finite-dimensional, then $T_t|_{H^-}$ is invertible for each $t \geq 0$. Therefore, we can set $T_{-t} := [T_t|_{H^-}]^{-1} : H^- \rightarrow H^-$ for each $t \geq 0$.

Consider the following semilinear stochastic evolution equation on H :

$$\begin{aligned} du(t) &= [-Au(t) + F(u(t))] dt + B_0 dW(t), \quad t \geq 0, \\ u(0) &= x \in H. \end{aligned} \tag{4.1}$$

In the above equation, let $F : H \rightarrow H$ be a globally Lipschitz map with Lipschitz constant L :

$$|F(v_1) - F(v_2)| \leq L|v_1 - v_2|, \quad v_1, v_2 \in H.$$

Then (4.1) has a unique mild solution given by

$$u(t, x) = T_t x + \int_0^t T_{t-s} F(u(s, x)) ds + \int_0^t T_{t-s} B_0 dW(s), \quad t \geq 0 \tag{4.2}$$

Furthermore, if $F : H \rightarrow H$ is $C^{k, \epsilon}$, the mild solution of (4.2) generates a $C^{k, \epsilon}$ perfect cocycle also denoted by $u : \mathbf{R}^+ \times H \times \Omega \rightarrow H$.

Suppose that $F : H \rightarrow H$ is globally bounded, and its Lipschitz constant L satisfies

$$L[\mu_{m+1}^{-1} - \mu_m^{-1}] < 1. \tag{4.3}$$

Note that the above condition is automatically satisfied in the affine linear case $F \equiv 0$.

The next proposition is key to the existence and uniqueness of a stationary random point for the cocycle (u, θ) in the sense of Definition 1.1.

Proposition 4.1.

Assume the above conditions on A, B_0, F together with (4.3). Then there is a unique $\mathcal{F}$ -measurable map $Y : \Omega \rightarrow H$ satisfying

$$\begin{aligned} Y(\omega) = & \int_{-\infty}^0 T_{-s} p^+ F(Y(\theta(s, \omega))) ds - \int_0^{\infty} T_{-s} p^- F(Y(\theta(s, \omega))) ds \\ & + (\omega) \int_{-\infty}^0 T_{-s} p^+ B_0 dW(s) - (\omega) \int_0^{\infty} T_{-s} p^- B_0 dW(s) \end{aligned} \quad (4.4)$$

for all $\omega \in \Omega$.

Proof.

We use a contraction mapping argument to show that the integral equation (4.4) has an $\mathcal{F}$ -measurable solution $Y : \Omega \rightarrow H$.

Define the $\mathcal{F}$ -measurable map $Y_1 : \Omega \rightarrow H$ by

$$Y_1(\omega) := (\omega) \int_{-\infty}^0 T_{-s} p^+ B_0 dW(s) - (\omega) \int_0^{\infty} T_{-s} p^- B_0 dW(s), \quad \omega \in \Omega.$$

Denote by $B(\Omega, H)$ the Banach space of all (surely) bounded $\mathcal{F}$ -measurable maps $Z : \Omega \rightarrow H$ given the supremum norm $\|Z\|_{\infty} := \sup_{\omega \in \Omega} |Z(\omega)|$. Define the map $\mathcal{M} : B(\Omega, H) \rightarrow L^0(\Omega, H)$ by

$$\begin{aligned} \mathcal{M}(Z)(\omega) := & \int_{-\infty}^0 T_{-s} p^+ F(Z(\theta(s, \omega)) + Y_1(\theta(s, \omega))) ds \\ & - \int_0^{\infty} T_{-s} p^- F(Z(\theta(s, \omega)) + Y_1(\theta(s, \omega))) ds \end{aligned}$$

for all $Z \in B(\Omega, H)$ and all $\omega \in \Omega$.

Note first that $\mathcal{M}$ maps $B(\Omega, H)$ into itself. To see this let $Z \in B(\Omega, H)$ and $\omega \in \Omega$. Then

$$\begin{aligned} |\mathcal{M}(Z)(\omega)| & \leq \|F\|_{\infty} \left[\int_{-\infty}^0 \|T_{-s} p^+\| ds + \int_0^{\infty} \|T_{-s} p^-\| ds \right] \\ & \leq \|F\|_{\infty} \left[\int_{-\infty}^0 e^{s\mu_{m+1}} ds + \int_0^{\infty} e^{s\mu_m} ds \right] \\ & \leq \|F\|_{\infty} [\mu_{m+1}^{-1} - \mu_m^{-1}] < \infty \end{aligned}$$

where $\|F\|_{\infty} := \sup_{v \in H} |F(v)|$. Hence $\mathcal{M}(Z) \in B(\Omega, H)$ for all $Z \in B(\Omega, H)$.

Secondly, $\mathcal{M}$ is a contraction. To prove this, take any $Z_1, Z_2 \in B(\Omega, H)$ and $\omega \in \Omega$. Then from the definition of $\mathcal{M}$, we get

$$\begin{aligned}
|\mathcal{M}(Z_1)(\omega) - \mathcal{M}(Z_2)(\omega)| &\leq L \int_{-\infty}^0 \|T_{-s}p^+\| \cdot |Z_1(\theta(s, \omega)) - Z_2(\theta(s, \omega))| ds \\
&\quad + L \int_0^{\infty} \|T_{-s}p^-\| \cdot |Z_1(\theta(s, \omega)) - Z_2(\theta(s, \omega))| ds \\
&\leq L \|Z_1 - Z_2\|_{\infty} \left[\int_{-\infty}^0 \|T_{-s}p^+\| ds + \int_0^{\infty} \|T_{-s}p^-\| ds \right] \\
&\leq L \|Z_1 - Z_2\|_{\infty} \left[\int_{-\infty}^0 e^{s\mu_{m+1}} ds + \int_0^{\infty} e^{s\mu_m} ds \right] \\
&= L[\mu_{m+1}^{-1} - \mu_m^{-1}] \|Z_1 - Z_2\|_{\infty} \\
&= \mu \|Z_1 - Z_2\|_{\infty}
\end{aligned}$$

where $\mu := L[\mu_{m+1}^{-1} - \mu_m^{-1}] < 1$. This proves that $\mathcal{M} : B(\Omega, H) \rightarrow B(\Omega, H)$ is a contraction, and hence has a unique fixed point $Z_0 \in B(\Omega, H)$. That is

$$\begin{aligned}
Z_0(\omega) &:= \int_{-\infty}^0 T_{-s}p^+ F(Z_0(\theta(s, \omega)) + Y_1(\theta(s, \omega))) ds \\
&\quad - \int_0^{\infty} T_{-s}p^- F(Z_0(\theta(s, \omega)) + Y_1(\theta(s, \omega))) ds
\end{aligned}$$

for all $\omega \in \Omega$. Now define $Y : \Omega \rightarrow H$ by

$$Y(\omega) := Z_0(\omega) + Y_1(\omega), \quad \omega \in \Omega.$$

It is easy to check that Y satisfies the identity (4.4).

Since Z_0 is uniquely determined, then so is Y . $\square$

The following proposition gives existence and uniqueness of a stationary point for the see (4.1).

Proposition 4.2.

Assume all the conditions on A, B_0, F stated in Proposition 4.1. Suppose that F is globally bounded, globally Lipschitz and satisfies condition (4.3). Then the semilinear see (4.1) has a unique

stationary point $Y : \Omega \rightarrow H$, i.e. $u(t, Y(\omega), \omega) = Y(\theta(t, \omega))$ for all $t \geq 0$ and $\omega \in \Omega$. Furthermore, $Y \in L^p(\Omega, H)$ for all $p \geq 1$.

Proof.

By hypotheses and Proposition 4.1, the integral equation (4.4) has a unique $\mathcal{F}$ -measurable solution $Y : \Omega \rightarrow H$. Let $t \geq 0$. Using (4.4), it follows that

$$\begin{aligned}
Y(\theta(t, \omega)) &= \int_{-\infty}^0 T_{-s} p^+ F(Y(\theta(t+s, \omega))) ds - \int_0^\infty T_{-s} p^- F(Y(\theta(t+s, \omega))) ds \\
&\quad + (\omega) \int_{-\infty}^0 T_{-s} p^+ B_0 dW(s+t) - (\omega) \int_0^\infty T_{-s} p^- B_0 dW(s+t) \\
&= \int_{-\infty}^t T_{t-s} p^+ F(Y(\theta(s, \omega))) ds - \int_t^\infty T_{t-s} p^- F(Y(\theta(s, \omega))) ds \\
&\quad + (\omega) \int_{-\infty}^t T_{t-s} p^+ B_0 dW(s) - (\omega) \int_t^\infty T_{t-s} p^- B_0 dW(s) \\
&= T_t \left[\int_{-\infty}^0 T_{-s} p^+ F(Y(\theta(s, \omega))) ds - \int_0^\infty T_{-s} p^- F(Y(\theta(s, \omega))) ds \right. \\
&\quad \left. + (\omega) \int_{-\infty}^0 T_{-s} p^+ B_0 dW(s) - (\omega) \int_0^\infty T_{-s} p^- B_0 dW(s) \right] \\
&\quad + \int_0^t T_{t-s} p^+ F(Y(\theta(s, \omega))) ds + \int_0^t T_{t-s} p^- F(Y(\theta(s, \omega))) ds \\
&\quad + (\omega) \int_0^t T_{t-s} p^+ B_0 dW(s) + (\omega) \int_0^t T_{t-s} p^- B_0 dW(s) \\
&= T_t Y(\omega) + \int_0^t T_{t-s} F(Y(\theta(s, \omega))) ds + (\omega) \int_0^t T_{t-s} B_0 dW(s).
\end{aligned}$$

This gives

$$Y(\theta(t, \omega)) = T_t Y(\omega) + \int_0^t T_{t-s} F(Y(\theta(s, \omega))) ds + (\omega) \int_0^t T_{t-s} B_0 dW(s)$$

for all $t \geq 0$. Therefore, $Y(\theta(t, \omega)), t \geq 0, \omega \in \Omega$, is a stationary solution of (4.2) (with $x = Y(\omega)$).

Since $u(t, Y(\omega), \omega), t \geq 0, \omega \in \Omega$, is also a solution of (4.2), then by uniqueness of the solution to (4.2), we must have

$$u(t, Y(\omega), \omega) = Y(\theta(t, \omega))$$

for all $t \geq 0$ and all $\omega \in \Omega$. Hence Y is a stationary point for the see (4.1).

The stationary point for (4.1) is unique (within the class of $\mathcal{F}$ -measurable maps $\Omega \rightarrow H$). To see this, it is sufficient to observe that the above computation shows that every stationary point of (4.1) is a solution of the integral equation (4.4). Uniqueness of the stationary solution then follows from Proposition 4.1.

In view of the proof of Proposition 4.1, the last assertion of Proposition 4.2 follows from the fact that $Y_1 \in L^p(\Omega, H)$ for all $p \geq 1$ and $Z_0 \in L^\infty(\Omega, H)$. $\square$.

The existence of local stable and unstable manifolds near a stationary point of the affine stochastic evolution equation (4.1) follows from a straightforward modification of the proof of Theorem 4.1 in the next section.

(b) Semilinear stochastic evolution equations: Linear noise

Here we recall the setting and hypotheses leading to Theorem 2.6 in [M-Z-Z.1].

We will prove the existence of local stable and unstable manifolds for semiflows generated by mild solutions of semilinear stochastic evolution equations of the form:

$$\left. \begin{aligned} du(t) &= -Au(t)dt + F(u(t))dt + Bu(t)dW(t), \quad t > 0, \\ u(0) &= x \in H. \end{aligned} \right\} \quad (4.5)$$

In the above equation $A : D(A) \subset H \rightarrow H$ is a closed linear operator on a separable real Hilbert space H . Assume that A has a complete orthonormal system of eigenvectors $\{e_n : n \geq 1\}$ with corresponding positive eigenvalues $\{\mu_n, n \geq 1\}$; i.e., $Ae_n = \mu_n e_n, n \geq 1$. Suppose $-A$ generates a strongly continuous semigroup of bounded linear operators $T_t : H \rightarrow H, t \geq 0$. Let E be a separable real Hilbert space. Suppose $W(t), t \geq 0$, is E -valued cylindrical Brownian motion defined on the canonical filtered Wiener space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, P)$ and with a separable covariance Hilbert space K , where $K \subset E$ is a Hilbert-Schmidt embedding. That is, Ω is the space of all continuous paths $\omega : \mathbf{R} \rightarrow E$ such that $\omega(0) = 0$ with the compact open topology, $\mathcal{F}$ is its Borel σ -field, $\mathcal{F}_t$ is the sub- σ -field generated by all evaluations $\Omega \ni \omega \mapsto \omega(u) \in E, u \leq t$, and P is Wiener measure on Ω . The Brownian motion is given by

$$W(t, \omega) := \omega(t), \quad \omega \in \Omega, t \in \mathbf{R},$$

and may be represented by

$$W(t) = \sum_{k=1}^{\infty} W^k(t) f_k, \quad t \in \mathbf{R},$$

where $\{f_k : k \geq 1\}$ is a complete orthonormal basis of K , and the $W^k, k \geq 1$, are standard independent one-dimensional Wiener processes ([D-Z.1], Chapter 4).

Suppose $B : H \rightarrow L_2(K, H)$ is a bounded linear operator. The stochastic integral in (4.5) is defined in the sense of ([D-Z.1], Chapter 4).

Assume the hypotheses of Theorem 2.4 in [M-Z-Z.1].

We will denote by $\theta : \mathbf{R} \times \Omega \rightarrow \Omega$ the standard P -preserving ergodic Wiener shift on Ω :

$$\theta(t, \omega)(s) := \omega(t + s) - \omega(t), \quad t, s \in \mathbf{R}.$$

Let $L(H)$ be the Banach space of all bounded linear operators $H \rightarrow H$ given the uniform operator norm $\|\cdot\|$. Denote by $L_2(H) \subset L(H)$ the Hilbert space of all Hilbert-Schmidt operators $S : H \rightarrow H$.

Suppose $F : H \rightarrow H$ is a (Fréchet) $C^{k, \epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) non-linear map satisfying the following Lipschitz and linear growth hypotheses:

$$\left. \begin{aligned} |F(v)| &\leq C(1 + |v|), \quad v \in H \\ |F(v_1) - F(v_2)| &\leq L_n |v_1 - v_2|, \quad v_i \in H, |v_i| \leq n, i = 1, 2, \end{aligned} \right\} \quad (4.6)$$

for some positive constants $C, L_n, n \geq 1$.

The mild solutions of the see (4.5) generate a $C^{k, \epsilon}$ ($k \geq 1, \epsilon \in (0, 1]$) perfect cocycle (U, θ) on H , satisfying all the assertions of Theorem 2.6 of [M-Z-Z.1].

Under the above conditions, one gets the following stable manifold theorem for hyperbolic stationary trajectories of the see (4.5).

Theorem 4.1.

Assume the above hypotheses on the coefficients of the see (4.5). Assume that the stochastic semiflow $U : \mathbf{R}^+ \times H \times \Omega \rightarrow H$ generated by mild solutions of (4.5) has a hyperbolic stationary point $Y : \Omega \rightarrow H$ such that $E \log^+ |Y| < \infty$. Then (U, θ) has a perfect family of $C^{k, \epsilon}$ local stable and unstable manifolds satisfying all the assertions of Theorem 2.1.

Proof.

One first checks the estimate

$$\int_{\Omega} \log^+ \sup_{0 \leq t_1, t_2 \leq a} \|U(t_2, Y(\theta(t_1, \omega)) + (\cdot), \theta(t_1, \omega))\|_{k, \epsilon} dP(\omega) < \infty \quad (4.7)$$

for any fixed $0 < \rho, a < \infty, k \geq 1$ and $\epsilon \in (0, 1]$. This estimate follows from the integrability condition on Y and assertion (vi) of Theorem 2.6 in [M-Z-Z.1]. The conclusion of Theorem 4.1 now follows immediately from Theorem 2.1. $\square$

(c) *Semilinear parabolic spde's: Lipschitz nonlinearity*

Consider the Laplacian

$$\Delta := \frac{1}{2} \sum_{i,j=1}^d \frac{\partial^2}{\partial \xi_i^2} \quad (4.8)$$

defined on a smooth bounded domain $\mathcal{D}$ in $\mathbf{R}^d$, with a smooth boundary $\partial\mathcal{D}$ with zero Dirichlet boundary conditions. Assume that $f : \mathbf{R} \rightarrow \mathbf{R}$ is a C_b^∞ function and let $d\xi$ be Lebesgue measure on $\mathbf{R}^d$. Let $W_n, n \geq 1$, be independent one-dimensional standard Brownian motions with $W_n(0) = 0$ defined on the canonical filtered Wiener space $(\Omega, \mathcal{F}, P, (\mathcal{F}_t)_{t \in \mathbf{R}})$. Let θ denote the Brownian shift on $\Omega := C(\mathbf{R}, \mathbf{R}^\infty)$. Recall that the Sobolev space $H_0^k(\mathcal{D})$ is the completion of $C_0^\infty(\mathcal{D}, \mathbf{R})$ under the Sobolev norm

$$\|u\|_{H_0^k(\mathcal{D})}^2 := \sum_{|\alpha| \leq k} \int_{\mathcal{D}} |D^\alpha u(\xi)|^2 d\xi.$$

Suppose further that $\sigma_n \in H_0^s(\mathcal{D}), n \geq 1$, and the series $\sum_{n=1}^\infty \sigma_n$ converges absolutely in $H_0^s(\mathcal{D})$ where $s > k + \frac{d}{2} > d$.

By Theorem 3.5 ([M-Z-Z.1]), weak solutions of the initial-value problem:

$$\left. \begin{aligned} du(t) &= \frac{1}{2} \Delta u(t) dt + f(u(t)) dt + \sum_{n=1}^\infty \sigma_n(\xi) u(t) dW_n(t), \quad t > 0 \\ u(0) &= \psi \in H_0^k(\mathcal{D}) \end{aligned} \right\} \quad (4.9)$$

give a perfect smooth cocycle (U, θ) on the Sobolev space $H_0^k(\mathcal{D})$ which satisfies all the assertions of Theorem 3.5 in [M.Z.Z.1]. Applying Theorem 2.1, we get the following stable manifold theorem for the spde (4.9):

Theorem 4.2.

Assume the above hypotheses on the coefficients of the spde (4.9). Assume that the stochastic semiflow $U : \mathbf{R}^+ \times H_0^k(\mathcal{D}) \times \Omega \rightarrow H_0^k(\mathcal{D})$ generated by weak solutions of (4.9) has a hyperbolic stationary point $Y : \Omega \rightarrow H_0^k(\mathcal{D})$ such that $E \log^+ \|Y\|_{H_0^k} < \infty$. Then (U, θ) has a perfect family of C^∞ local stable and unstable manifolds in $H_0^k(\mathcal{D})$ satisfying all the assertions of Theorem 2.1.

(d) Stochastic reaction diffusion equations: dissipative nonlinearity

In section 4 (a) of [M.Z.Z.1], we constructed a C^1 stochastic semiflow on the Hilbert space $H := L^2(\mathcal{D})$ for a stochastic reaction-diffusion equation

$$du = \nu \Delta u dt + u(1 - |u|^\alpha) dt + \sum_{i=1}^{\infty} \sigma_i(\xi) u(t) dW_i(t), \quad (4.10)$$

defined on a bounded domain $\mathcal{D} \subset \mathbf{R}^d$ with a smooth boundary $\partial\mathcal{D}$. In (4.10), the Laplacian on $\mathcal{D}$ is denoted by Δ , and we impose Dirichlet boundary conditions on $\partial\mathcal{D}$. The $W_i, i \geq 1$, are independent one-dimensional standard Brownian motions and $\sum_{i=1}^{\infty} \sigma_i$ is absolutely convergent in $H^s(\mathcal{D})$, for $s > 2 + \frac{d}{2}$. The dissipative term yields the existence of a unique stationary solution of (4.10) under a suitable choice of the parameter ν ([D-Z.2]).

In view of the estimates in Theorem 4.1 ([M.Z.Z.1]) and Theorem 2.1, one gets the following:

Theorem 4.3.

Assume the above hypotheses on the coefficients of the spde (4.10). Let $\alpha < \frac{4}{d}$. Assume that the stochastic semiflow $U : \mathbf{R}^+ \times L^2(\mathcal{D}) \times \Omega \rightarrow L^2(\mathcal{D})$ generated by mild solutions of (4.10) has a hyperbolic stationary point $Y : \Omega \rightarrow L^2(\mathcal{D})$ such that $E \log^+ \|Y\|_{L^2} < \infty$. Then (U, θ) has a perfect family of C^1 local stable and unstable manifolds in $L^2(\mathcal{D})$ satisfying all the assertions of Theorem 2.1.

Remarks.

- (i) The results in Sections (c) and (d) hold if the Euclidean domain $\mathcal{D}$ is replaced by a compact smooth d -dimensional Riemannian manifold M (possibly with a smooth boundary ∂M).

- (ii) We conjecture that Theorem 4.3 still holds (but with *Lipschitz* stable/unstable manifolds) if the dissipative term $u(1 - |u|^\alpha)$ is replaced by a more general one of the form $F(u) := f \circ u$, where $f : \mathbf{R} \rightarrow \mathbf{R}$ is a C^1 function satisfying the following classical estimates:

$$-c_1 - \alpha_1|x|^p \leq f(x)x \leq c_1 - \alpha_2|x|^p, \quad f'(x) \leq c_2,$$

for all $x \in \mathbf{R}$, with $c_1, c_2, \alpha_1, \alpha_2$ positive constants, and p any integer greater than 2.

- (iii) Is it true that the stochastic flow and the local stable/unstable manifolds in Theorem 4.3 are of class C^2 ?

(e) *Stochastic Burgers equation: additive noise*

The existence of a C^1 stochastic semiflow on $L^2([0, 1])$ for Burgers equation

$$du + u \frac{\partial u}{\partial \xi} dt = \nu \Delta u dt + dW(t), \quad t > 0, \quad \nu > 0, \quad (4.11)$$

was established in Part I of this work, where $W(t)$, $t > 0$, is an infinite dimensional Brownian motion on $L^2[0, 1]$. See [M-Z-Z.1], Theorem 4.3.

Under extra spatial smoothness hypotheses on the noise, viz. $W(t, \cdot) \in C^3([0, 1])$, Burgers equation (4.11) admits a unique stationary point ([Si]). More generally, with our weaker condition on the noise W ([M.Z.Z.1], Section 4 (b)), we stipulate that equation (4.11) has a hyperbolic stationary point. In this case, we get the following result:

Theorem 4.4.

Assume the hypotheses of Theorem 4.3 of [M.Z.Z.1] on the coefficients of Burgers spde (4.11). Assume that the stochastic semiflow $U : \mathbf{R}^+ \times L^2([0, 1]) \times \Omega \rightarrow L^2([0, 1])$ generated by mild solutions of (4.11) has a hyperbolic stationary point $Y : \Omega \rightarrow L^2([0, 1])$ such that $E \log^+ \|Y\|_{L^2} < \infty$. Then (U, θ) has a perfect family of C^1 local stable and unstable manifolds in $L^2([0, 1])$ satisfying all the assertions of Theorem 2.1.

Note that hyperbolicity of the stationary point in Theorem 4.4 is in the sense of Definition (1.3). Theorem 1.1 of this article and Theorem 4.3 of ([M.Z.Z.1]) imply that the Lyapunov spectrum

for the linearization of (4.11) exists and is discrete for any viscosity $\nu > 0$. When W is C^3 in the space variable, it is known that for any C^2 initial condition, the solution $u(t)$ of (4.11) converges to the stationary solution for any positive viscosity $\nu > 0$ ([Si]). It is therefore easy to see that the stable manifold is the whole of $L^2[0, 1]$.

The case of sufficiently large viscosity and rough noise $W(t) \in L^2([0, 1])$ is currently being studied ([L-Z]). This work shows that (4.11) admits a unique globally exponentially stable stationary point in this case. So in this (somewhat non-generic) case, the unstable manifold consists of the single random point $Y(\omega) \in L^2([0, 1])$, and the non-linear cocycle U will approach $Y(\omega)$ with exponential speed less than or equal to the top Lyapunov exponent λ_1 of the linearized Burgers equation.

We conjecture that the assertions in the above paragraph still hold for any viscosity $\nu > 0$ (cf. [D-Z.2], Theorem 14.4.4). Further analysis of the Lyapunov spectrum for (4.11) (in the cases of small and zero viscosity ν) is postponed to a future project.

Acknowledgments

The authors would like to thank Prof. B. Øksendal for inviting them to Oslo in the summer 2000 where the project was started. They are also grateful to Prof. K.D. Elworthy for inviting them to Warwick on numerous occasions especially during the Warwick SPDE's Symposium 2000/2001 so that they may have opportunities to meet; to Prof. A. Truman for inviting them to the International Workshop of Probabilistic Methods in Fluids at Swansea in April 2002 where preliminary versions of the results were announced; and to the organizers of ICM 2002 and First Sino-German Stochastic Analysis Conference in Beijing in August 2002 where the results were presented. We would also like to thank Z. Brzezniak, F. Flandoli, Y. Liu, J. Robinson, J. Zabczyk, and several other colleagues for useful discussions.

REFERENCES

- [Co] Cohn, D. L., *Measure Theory*, Birkhäuser (1980).
- [C-V] Castaing, C., and Valadier, M., *Convex Analysis and Measurable Multifunctions*, Lecture Notes in Mathematics, vol. 580, Springer-Verlag, Berlin-Heidelberg-New York (1977).
- [D-Z.1] Da Prato, G., and Zabczyk, J., *Stochastic Equations in Infinite Dimensions*, Cambridge University Press (1992).
- [D-Z.2] Da Prato, G., and Zabczyk, J., *Ergodicity for Infinite Dimensional Systems*, Cambridge University Press (1996).
- [F-S] Flandoli, F., and Schaumlöffel, K.-U., A multiplicative ergodic theorem with applications to a first order stochastic hyperbolic equation in a bounded domain, *Stochastics and Stochastic Reports*, 34 (1991), no. 3-4, 241–255.
- [L-Z] Liu, Y. and Zhao, H.Z., The stationary solutions of some SPDEs, in preparation.
- [Mo.1] Mohammed, S.-E. A., The Lyapunov spectrum and stable manifolds for stochastic linear delay equations, *Stochastics and Stochastic Reports*, Vol. 29 (1990), 89-131.
- [M-S.1] Mohammed, S.-E. A., and Scheutzow, M. K. R., Lyapunov exponents of linear stochastic functional differential equations driven by semimartingales, Part I: The multiplicative ergodic theory, *Ann. Inst. Henri Poincaré, Probabilités et Statistiques*, Vol. 32, 1, (1996), 69-105. pp. 43.
- [M-S.2] Mohammed, S.-E. A., and Scheutzow, M. K. R., Spatial estimates for stochastic flows in Euclidean space, *The Annals of Probability*, 26, 1, (1998), 56-77.
- [M-S.3] Mohammed, S.-E. A., and Scheutzow, M. K. R., The stable manifold theorem for stochastic differential equations, *The Annals of Probability*, Vol. 27, No. 2, (1999), 615-652.
- [M-S.4] Mohammed, S.-E. A., and Scheutzow, M. K. R., The stable manifold theorem for non-linear stochastic systems with memory. Part I: Existence of the semiflow, Part II: The local stable manifold theorem. (preprints).
- [M-Z-Z.1] Mohammed, S.-E. A., Zhang, T. S. and Zhao, H., The stable manifold theorem for semilinear stochastic evolution equations and stochastic partial differential equations I: Flows and stationary solutions (preprint) (2002).
- [O] Oseledec, V. I., A multiplicative ergodic theorem. Lyapunov characteristic numbers for dynamical systems, *Trudy Moskov. Mat. Obšč.* 19 (1968), 179-210. English transl. *Trans. Moscow Math. Soc.* 19 (1968), 197-221.
- [Ru.1] Ruelle, D., Ergodic theory of differentiable dynamical systems, *Publ. Math. Inst. Hautes Etud. Sci.* (1979), 275-306.
- [Ru.2] Ruelle, D., Characteristic exponents and invariant manifolds in Hilbert space, *Annals of Mathematics* 115 (1982), 243–290.

- [Si] Sinai, Ya. G., Burgers system driven by a periodic stochastic flow, In: *Itô's stochastic calculus and probability theory*, Springer, Tokyo (1996), 347–353.

Salah-Eldin A. Mohammed
Department of Mathematics,
Southern Illinois University at Carbondale,
Carbondale, Illinois 62901.
Email: salah@sfde.math.siu.edu
Web page: <http://sfde.math.siu.edu>

Tusheng Zhang
Department of Mathematics
University of Manchester,
Oxford Road, Manchester M13 9PL
UK.
Email: tzhang@math.man.ac.uk

Huaizhong Zhao
Department of Mathematical Sciences
Loughborough University,
LE11 3TU,
UK.
Email: H.Zhao@lboro.ac.uk